

# The Physics of Neutrino Interactions

**M. Sajjad Athar**  
**S. K. Singh**



# Contents

<i>List of Figures</i>	<i>xxi</i>
<i>List of Tables</i>	<i>xxxv</i>
<i>Preface</i>	<i>xxxix</i>
<i>Acknowledgments</i>	<i>xliv</i>
<b>1 Neutrino Properties and Its Interactions</b>	<b>1</b>
1.1 Historical Introduction to Neutrinos	1
1.1.1 Neutrino hypothesis	1
1.1.2 The problem of energy conservation in $\beta$ -decays of nuclei	2
1.1.3 Anomalies in the spin–statistics relation for nuclei	3
1.1.4 Pauli’s neutron/neutrino vs. Fermi’s neutrino	4
1.2 Neutrino Interactions	5
1.2.1 Fermi theory of $\beta$ -decay	6
1.2.2 Gamow–Teller theory	8
1.2.3 Parity violation and the two-component neutrino	10
1.2.4 Chiral ( $\gamma_5$ ) invariance and $V - A$ theory of $\beta$ -decays	11
1.2.5 Intermediate vector boson (IVB)	12
1.2.6 Weak interactions in strangeness sector: The Cabibbo theory, and the GIM mechanism	14
1.2.7 Quark flavors and the CKM matrix	16
1.2.8 Nonleptonic weak interaction and CP violation	17
1.3 Neutrino Flavors and Universality of Neutrino Interactions	18
1.3.1 Experimental discovery of $\bar{\nu}_e$ and $\bar{\nu}_e \neq \nu_e$	18
1.3.2 Discovery of muons and muon neutrinos	20
1.3.3 Lepton number conservation and $e - \mu$ universality	21
1.3.4 Discovery of tau neutrino and $e - \mu - \tau$ universality	22
1.4 Properties of Neutrinos	23
1.4.1 Weyl, Dirac and Majorana neutrinos	23
1.4.2 Neutrino mass	26
1.4.3 Neutrino charge and charge radius	28

4.4	Dyson Expansion and Wick's Theorem	146
4.4.1	Dyson expansion	146
4.4.2	Wick's theorem	148
4.5	S-matrix and Feynman Diagrams	151
4.6	Invariant Matrix Elements and Feynman Diagrams	156
4.6.1	Matrix elements in first order perturbation theory	156
4.6.2	Matrix elements in second order perturbation theory	158
4.6.3	Matrix elements for the closed loops	160
4.6.4	Feynman rules: A summary	161
4.7	Scattering Cross Sections and Particle Decay Rates	163
4.7.1	Scattering cross sections	163
4.7.2	Particle decay rates	165
<b>5</b>	<b>Phenomenological Theory I: Nuclear <math>\beta</math>-decays and Weak Interaction of Leptons</b>	<b>167</b>
5.1	Introduction	167
5.2	Development of Phenomenological Theory	168
5.2.1	Fermi and Gamow–Teller theories of $\beta$ -decays	168
5.2.2	General form of the Hamiltonian and the parity violation in $\beta$ -decays	177
5.2.3	The energy and angular distribution of electrons (positrons) for the $\beta^+$ ( $\beta^-$ )-decay of unpolarized nuclei	180
5.2.4	The longitudinal polarization of $e^-$ ( $e^+$ ) from $\beta^-$ ( $\beta^+$ )-decays of unpolarized nuclei	182
5.2.5	Helicity of the neutrino	183
5.2.6	Spin–momentum correlations in the $\beta$ -decay of polarized nuclei	186
5.2.7	Mixed $\beta$ -transitions and sign of $\frac{C_A}{C_V}$	186
5.3	Two-component Neutrino and the $V - A$ Theory	188
5.4	Weak Interaction of Muon	189
5.4.1	Weak decay of muons	190
5.4.2	General structure of weak interaction in muon-decay and Michel parameters	196
5.4.3	Radiative corrections to $\mu$ -decays	198
5.5	Inverse Muon-decay and $\nu_\mu$ Scattering	200
5.6	Muon Capture and $\mu - e$ Universality	202
5.7	Limitations of the Phenomenological Theory	203
5.7.1	High energy behavior of $\nu_l - l^-$ scattering and unitarity	203
5.7.2	Divergence and renormalization	204
5.7.3	Intermediate vector boson (IVB) theory	205
5.7.4	Radiative corrections	208
5.8	$\tau$ Lepton and Its Weak Decays and $e - \mu - \tau$ Universality	209
5.8.1	$\tau$ lepton and its properties	209

5.8.2 Weak decays of $\tau$ leptons and $e - \mu - \tau$ - universality	212
<b>6 Phenomenological Theory II: Weak Decays of Hadrons</b>	<b>215</b>
6.1 Introduction	215
6.2 Semileptonic Weak Decays of Hadrons without Strangeness	217
6.2.1 Two-body decay of pions: $\pi l_2$ decays	217
6.2.2 Three-body decays of pions: $\pi l_3$ decays	221
6.3 Symmetry Properties of the Weak Hadronic Current	222
6.3.1 Lorentz transformation properties and matrix elements	223
6.3.2 Isospin properties of the weak hadronic current	224
6.3.3 T invariance	225
6.3.4 Conserved vector current hypothesis	226
6.3.5 Implications of the CVC hypothesis	226
6.3.6 Partial conservation of axial vector current (PCAC)	228
6.3.7 Implications of PCAC	229
6.3.8 G-parity and second class currents	233
6.4 Semileptonic Weak Decays of Hadrons with Strangeness	234
6.4.1 The Cabibbo theory and the universality of weak interactions	236
6.4.2 The Cabibbo theory in the quark model and quark mixing	237
6.4.3 Applications of Cabibbo theory: K decays	239
6.4.4 Semileptonic decays of hyperons (Y)	242
6.4.5 Physical observables in semileptonic hyperon decays	244
6.5 Nonleptonic Decays of Strange Particles	247
6.5.1 Nonleptonic decays of K-mesons	247
6.5.2 Nonleptonic decays of hyperons	249
6.5.3 Radiative weak decays	251
6.6 CP Violation in the Neutral Kaon Sector	253
6.6.1 Neutral kaons, CP eigenstates, and $K^0 - \bar{K}^0$ oscillations	253
6.6.2 CP violation in the neutral kaon decays	257
6.7 Flavour Changing Neutral Currents (FCNC) and GIM Mechanism	259
6.7.1 Six quark mixing and CKM matrix	262
6.8 Weak Interaction of Hadrons with Charm and Heavy Flavors	264
6.8.1 Discovery of charm and heavy flavors	264
6.8.2 Weak decays of particles with charm and heavy flavors	266
6.8.3 Weak decays of particles with charm	266
6.8.4 Weak decays of particles with heavy flavors	269
6.9 Limitations of the Phenomenological Theory from the Hadron Sector	269

<b>7 Gauge Field Theories and Fundamental Interactions</b>	<b>274</b>
7.1 Introduction	274
7.2 Gauge Invariance in Field Theory	278
7.2.1 Maxwell's equations from a variational principle	279
7.2.2 Gauge invariance in field theory and phase invariance	280
7.3 Local Gauge Symmetries and Fundamental Interactions	282
7.3.1 Introduction	282
7.3.2 $U(1)$ gauge symmetry and the electromagnetic interactions of spin 0 particles	284
7.3.3 $U(1)$ gauge symmetry and the electromagnetic interactions of spin $\frac{1}{2}$ particles	287
7.3.4 $SU(2)$ gauge symmetry and Yang–Mills fields	289
7.3.5 $SU(3)_c$ gauge symmetry and QCD	294
7.3.6 Exact, broken, and spontaneously broken symmetries	296
7.3.7 Spontaneous symmetry breaking: Some formal results	299
7.3.8 Spontaneous symmetry breaking (SSB) and Goldstone's theorem	303
7.3.9 Spontaneously broken discrete symmetry: A simple model	307
7.3.10 Spontaneously broken continuous global symmetry under $U(1)$ gauge transformation and Goldstone boson	310
7.3.11 Spontaneously broken continuous local $U(1)$ symmetry and Higgs mechanism	313
7.3.12 Spontaneously broken local $SU(2)$ gauge symmetry and Higgs mechanism	315
<b>8 Unified Theory of Electroweak Interactions</b>	<b>318</b>
8.1 Introduction	318
8.2 Description of the Weinberg–Salam Model for Leptons	320
8.3 Predictions of the Weinberg–Salam Model	326
8.3.1 Masses of gauge bosons	326
8.3.2 Charged current weak interactions and electromagnetic interactions	328
8.3.3 Neutral current interaction and weak mixing angle	329
8.3.4 Mass of the electron	331
8.4 Extension to the Leptons of Other Flavors	331
8.4.1 Extension to the quark sector using GIM mechanism	335
8.4.2 Triumphs of the Weinberg–Salam–Glashow Model	337
8.4.3 Discovery of the neutral currents in $\nu$ -scattering	338
8.5 Discovery of Neutral Currents in Electron Scattering	341
8.6 Discovery of $W^\pm$ and Z Bosons	350
8.6.1 Properties of $W^\pm$ and Z boson	350
8.7 Higgs Boson	355
8.7.1 Discovery of Higgs boson	358

<b>9 Neutrino and Electron Scattering from Point Particles</b>	<b>362</b>
9.1 Introduction	362
9.2 $e^- + \mu^- \rightarrow e^- + \mu^-$ scattering	363
9.3 $\nu_\mu + e^- \rightarrow \mu^- + \nu_e$ scattering	368
9.4 $\nu_\mu + e^- \rightarrow \nu_\mu + e^-$ scattering	370
9.5 $\nu_e + e^- \rightarrow \nu_e + e^-$ scattering	373
9.5.1 Determination of the magnetic moment of neutrinos	377
9.6 $e^- + e^+ \rightarrow \mu^- + \mu^+$	378
<b>10 Neutrino scattering Cross Sections from Hadrons: Quasielastic Scattering</b>	<b>385</b>
10.1 Introduction	385
10.2 Physical Significance of the Form Factor	386
10.3 $e^- - \pi^\pm$ Elastic Scattering	389
10.4 Electromagnetic Scattering of Electrons with Nucleons	391
10.4.1 Matrix element and form factors	391
10.4.2 Physical interpretation of the form factors	394
10.4.3 Cross sections and the Rosenbluth separation	396
10.4.4 Experimental determination of the form factor	398
10.4.5 Numerical parameterization of the electromagnetic form factors	400
10.5 Quasielastic and Elastic $\nu$ Scattering Processes on Nucleons	402
10.5.1 Introduction	402
10.5.2 Interaction Lagrangian	403
10.5.3 Charged current quasielastic reaction and weak nucleon form factors	404
10.5.4 Neutral current elastic reactions and weak nucleon form factors	407
10.5.5 Symmetry properties of weak hadronic currents and form factors	408
10.5.6 Parameterization of the weak form factors	408
10.5.7 Cross sections for charged current processes	412
10.6 Quasielastic Hyperon Production	414
10.6.1 Matrix elements and form factors	414
10.6.2 Vector form factors	416
10.6.3 Axial vector form factors	416
10.6.4 Cross sections: Experimental results	417
10.7 Polarization of Final Hadrons and Leptons	418
10.7.1 Introduction	418
10.7.2 Polarization of the final hadron	419
10.7.3 Polarization of the final lepton	423

<b>11 Neutrino Scattering from Hadrons: Inelastic Scattering (I)</b>	<b>425</b>
11.1 Introduction	425
11.2 Inelastic Scattering through CC Excitation of Resonances	430
11.2.1 CC excitation of spin 1/2 resonances	430
11.2.2 CC excitation of spin 3/2 resonances	446
11.3 Neutral Current Reactions	457
11.3.1 Excitation of spin 1/2 resonances	457
11.3.2 Excitation of spin 3/2 resonances	461
11.4 Non-resonant Contributions	463
11.4.1 Chiral symmetry	465
11.4.2 Transformation of mesons under chiral transformation	466
11.4.3 Linear sigma model	468
11.4.4 Explicitly broken chiral symmetry	469
11.4.5 Non-linear sigma model	470
11.4.6 Lagrangian for the meson–meson and meson–gauge boson interactions	472
11.4.7 Lagrangian for the meson–baryon–gauge boson interaction	475
<b>12 Neutrino Scattering from Hadrons: Inelastic Scattering (II)</b>	<b>478</b>
12.1 Introduction	478
12.2 Single Pion Production	479
12.2.1 Charged current	480
12.2.2 Neutral current	485
12.2.3 Cross sections	487
12.3 Eta Production	490
12.3.1 Charged current	491
12.3.2 Neutral current	492
12.3.3 Cross section	493
12.4 Associated Production of Strange Particles	493
12.4.1 Charged current	495
12.4.2 Cross section	497
12.5 Kaon Production	498
12.5.1 Cross section	500
12.6 Antikaon Production	501
12.6.1 Cross section	505
<b>13 Neutrino Scattering from Hadrons: Deep Inelastic Scattering</b>	<b>507</b>
13.1 Introduction	507
13.2 Charged Lepton–nucleon DIS	511

13.2.1 Bjorken scaling and parton model	515
13.2.2 Differential scattering cross section in terms of dimensionless variables $x$ and $y$	525
13.3 Deep Inelastic Charged Current $\nu_l/\bar{\nu}_l - N$ Scattering	527
13.3.1 Relation between electromagnetic and weak structure functions	533
13.3.2 Experimental results of charged current total scattering cross section	534
13.4 Deep Inelastic Neutral Current $\nu_l/\bar{\nu}_l - N$ Scattering	536
13.5 QCD Corrections	539
13.5.1 Modified parton model	539
13.5.2 NLO evolution	542
13.5.3 TMC effect	544
13.5.4 Higher twist effect	547
13.6 Sum Rules in DIS	548
13.6.1 Adler sum rule	548
13.6.2 Gross–Llewellyn Smith sum rule	549
13.7 Quark–hadron (QH) Duality	550
13.8 Duality in Charged Lepton–nucleon Scattering	555
13.9 Duality in Neutrino–nucleon Scattering	558
<b>14 Weak Quasielastic <math>\nu(\bar{\nu})</math>-nucleus Scattering</b>	<b>561</b>
14.1 Introduction	561
14.2 Physics of Nuclear Medium Effects in Quasielastic Scattering	564
14.3 General Considerations	569
14.4 Low Energy Quasielastic Reactions	571
14.4.1 Multipole expansion of the matrix elements	571
14.4.2 Cross sections	575
14.4.3 Single particle matrix element in the shell model of nuclei	577
14.5 Quasielastic $\nu(\bar{\nu})$ Reactions at Intermediate Energies	578
14.5.1 Introduction	578
14.5.2 Fermi gas model	580
14.5.3 Local Fermi gas model	582
14.6 Cross Sections and Effect of Nuclear Medium	595
14.7 Nuclear Medium Effects in Neutrino Oscillation Experiments	599
<b>15 Inelastic Scattering of (Anti)neutrinos from Nuclei</b>	<b>603</b>
15.1 Introduction	603
15.2 Charged Current Inelastic Reactions	606
15.2.1 Incoherent meson production	606

15.2.2	Pion production in the delta dominance model	609
15.2.3	Quasielastic-like production of leptons	614
15.3	Coherent Pion Production	614
15.3.1	Introduction	614
15.3.2	PCAC based methods	616
15.4	Microscopic Model for Coherent Weak Pion Production	618
15.4.1	Cross sections	620
15.4.2	Final state interactions	620
15.5	Results for Cross Sections	623
15.6	Pion Production through Hyperon Excitation	627
15.6.1	Inelastic production of kaons	630
15.6.2	Inelastic production of photons	630
<b>16</b>	<b>Deep Inelastic Scattering of (Anti)neutrinos from Nuclei</b>	<b>632</b>
16.1	Introduction	632
16.2	DIS from Bound Nucleons	634
16.3	Extraction of Structure Functions from Cross Section Measurements	637
16.4	Phenomenological Study	639
16.5	Theoretical Study	645
16.5.1	Deep inelastic charged leptons and (anti)neutrino scattering from nuclei	645
16.5.2	Aligarh–Valencia model	647
16.5.3	Kulagin–Petti model	659
16.5.4	Isoscalarity corrections: Phenomenological approach	661
16.6	Results and Discussions	662
<b>17</b>	<b>Neutrino Sources and Detection of Neutrinos</b>	<b>669</b>
17.1	Introduction	669
17.2	Solar Neutrinos	670
17.2.1	Production	670
17.2.2	Detection	673
17.3	Atmospheric Neutrinos	677
17.3.1	Introduction	677
17.3.2	Detection	682
17.4	Reactor Antineutrinos	685
17.4.1	Detection	686
17.5	Supernova Neutrinos	687
17.5.1	Introduction	687
17.5.2	Neutrino emission in supernova explosions	691

17.5.3	Detection	692
17.6	Geoneutrinos	692
17.7	Relic Neutrinos	694
17.8	Accelerator Neutrinos	695
17.8.1	Wide and narrow band of neutrino beams	697
17.9	Neutrinos from the Decay at Rest (DAR) Sources	702
17.10	Neutrinos from Spallation Neutrons	703
17.11	Neutrinos from Muon Storage Ring (MSR)	704
17.12	Beta-beam Neutrinos	704
17.13	Very High Energy Cosmic Neutrinos	706
<b>18</b>	<b>Neutrino Mixing and Oscillations</b>	<b>708</b>
18.1	Introduction	708
18.2	Neutrino Mixing	710
18.2.1	Two-flavor neutrino oscillations in vacuum	710
18.2.2	Three-flavor neutrino oscillation in vacuum	718
18.2.3	Survival probability for $\nu_\alpha \rightarrow \nu_\alpha$	721
18.2.4	Transition probability for $\nu_\alpha \rightarrow \nu_\beta$	722
18.2.5	CP violation in the leptonic sector	723
18.2.6	Series expansions for neutrino oscillation probabilities	726
18.2.7	Neutrino mass hierarchy: Normal and inverted	729
18.3	Neutrino Oscillation in Matter	732
18.3.1	Effective potential for neutrino–matter interactions	733
18.3.2	Interaction Hamiltonian in matter	736
18.3.3	Probability for oscillation in matter	739
18.3.4	Resonance condition and level crossing	740
18.4	Neutrino Oscillation: Experimental Status	741
18.5	Sterile Neutrinos	743
18.6	Phenomenology of Sterile Neutrinos	745
18.7	Present and Future Experiments	746
<b>19</b>	<b>Neutrino Astrophysics and the Synthesis of Elements</b>	<b>749</b>
19.1	Introduction	749
19.2	The Big Bang and the Nucleosynthesis of the Lighter Elements	753
19.3	Interstellar Matter	756
19.4	Formation of Stars	757
19.4.1	The Hertzsprung–Russell diagram	758
19.5	Stellar Nucleosynthesis	760

19.5.1	Death of a star	760
19.5.2	Chandrasekhar limit	763
19.5.3	Death of middle mass stars	763
19.6	The Supernova Nucleosynthesis: Formation of Heavy Elements	767
19.6.1	Nucleosynthesis of neutron-rich elements	768
19.6.2	Nucleosynthesis of proton-rich elements	772
<b>20</b>	<b>Neutrino Interactions Beyond the Standard Model</b>	<b>775</b>
20.1	Introduction	775
20.2	Neutrinoless Double-beta Decay	776
20.2.1	General considerations	776
20.2.2	Decay rates of $0\nu\beta\beta$	779
20.2.3	Experiments	785
20.3	Lepton Flavor Violating Processes	788
20.4	Flavor Changing Neutral Currents	789
20.4.1	Particle decay processes	789
20.4.2	Production of strange and heavy flavored hadrons in scattering experiments	790
20.5	Nonstandard Interaction (NSI) in High Precision Low Energy Weak Processes	791
20.6	Summary	794
 <i>Appendices</i>		
<b>A</b>	<b>Lorentz Transformation and Covariance of the Dirac Equation</b>	<b>795</b>
A.1	Lorentz Transformations	795
A.2	Covariance of Dirac Equation	799
A.3	Bilinear Covariants	803
A.4	Nonrelativistic Reduction	804
<b>B</b>	<b>Cabibbo Theory</b>	<b>807</b>
B.1	Cabibbo Theory, SU(3) Symmetry, and Weak $N\text{--}Y$ Transition Form Factors	807
B.2	Octet Representation of Mesons	814
B.3	Octet Representation of Baryons	816
<b>C</b>	<b>Some Properties of Pauli and Dirac Matrices and Spin Density Matrices</b>	<b>819</b>
C.1	Trace Properties of Pauli and Dirac Matrices	819
C.2	Spin Density Matrix	821

---

<b>D Leptonic and Hadronic Tensors</b>	<b>828</b>
D.1 Contraction of Leptonic Tensors in Electromagnetic Interactions	828
D.2 Contraction of Leptonic Tensors in the Case of Weak Interactions	830
D.3 Contraction of Weak Leptonic Tensor with Hadronic Tensor	831
<b>E General Expression for the Total Scattering Cross Section and Decay Rates</b>	<b>833</b>
E.1 Cross Section	833
E.2 Decay Rate	837
<b>F Expressions of <math>N(q^2)</math>, the Coefficients of the Polarization Observables</b>	<b>841</b>
F.1 Expression of $N(q^2)$ in Terms of Mandelstam Variables	841
F.2 Expressions of $A^h(q^2)$ , $B^h(q^2)$ , and $C^h(q^2)$	842
F.3 Expressions of $A^l(q^2)$ , $B^l(q^2)$ and $C^l(q^2)$	844
<i>References</i>	847
<i>Index</i>	923

# Figures

1.1	Continuous $\beta$ -decay spectrum of RaE.	2
1.2	(a) Four fermions interact for $\beta$ -decay; (b) Interaction of two electrons with a photon field (QED).	7
1.3	$\beta$ -decay spectrum for GT transitions vs. momentum $ \vec{p}_e $ .	9
1.4	$\beta^-$ -decay spectrum with $m_\nu \neq 0$ and $m_\nu = 0$ .	11
1.5	$\beta^-$ -decay process mediated through a $W^-$ boson.	13
1.6	Divergence of cross section at high energies ( $\sigma \propto s$ ).	13
1.7	Interaction of quark with $W$ boson field.	15
1.8	Two instant and one delayed photon in Reines and Cowan's experiment.	19
1.9	Weyl neutrinos.	24
1.10	Dirac neutrinos.	24
1.11	Majorana neutrinos.	25
1.12	Neutrinoless double beta decay ( $0\nu\beta\beta$ ).	25
1.13	(a) $p\bar{p}$ collisions inside the CMS detector at CERN. This event was recorded in 2012 by the compact muon solenoid (CMS) at the large hadron collider (LHC). (b) Experimental confirmation of the Higgs boson.	34
1.14	A $W^+$ -boson interacts at the point $A$ and produces $e^+$ and $\nu_e$ . The $\nu_e$ travels a long distance; it interacts with the detector $B$ and produces a lepton $e^-$ as shown in the top panel.	37
1.15	Neutrino mass hierarchy.	40
2.1	Energy spectrum of Dirac particles with energy gap $2E_0$ .	61
2.2	Mirror reflection resulting $\vec{r} \rightarrow -\vec{r}$ while $ \vec{r} $ remains invariant.	81
2.3	Time reversal $t \rightarrow -t$ .	89
3.1	$\delta$ variation-extremum path.	98
3.2	Contours for Eq. (3.144). $C^+$ is for $\Delta_F^+$ and $C^-$ is for $\Delta_F^-$ .	119

3.3 Propagation of bosons between $x_1$ and $x_2$ for (a) $t_1 > t_2$ and (b) $t_2 > t_1$ .	121
3.4 Contour for the boson propagator $\Delta_F$ .	121
3.5 Position of displaced poles in the contour integration for $\Delta_F$ .	122
3.6 Propagation of a fermion between $x_1$ and $x_2$ for (a) $t_1 > t_2$ and (b) $t_2 > t_1$ .	130
3.7 Interaction of two electromagnetic currents through photon exchange.	138
4.1 $(t_1, t_2)$ plane for integration.	147
4.2 Feynman diagrams for eight basic processes in QED. (a) $e^+e^-$ annihilation, (b) $e^+$ scattering, (c) $e^-$ scattering, and (d) $e^+e^-$ creation, accompanied by the absorption (upper row) and emission (lower row) of a photon.	151
4.3 Feynman diagrams for (a) Compton scattering for $e^-$ , (b) Compton scattering for $e^+$ , (c) pair annihilation, and (d) pair creation.	153
4.4 Feynman diagrams for (a, b) Møller scattering of $e^- - e^-$ , (c, d) Møller scattering of $e^+ - e^+$ , and (e, f) Bhabha scattering of $e^- - e^+$ .	154
4.5 Electron (left) and positron (right) self energy.	155
4.6 Photon self energy.	155
4.7 The vacuum diagram.	156
4.8 The process $e^- \rightarrow e^- + \gamma$ .	158
4.9 Compton scattering of electrons via (a) s-channel and (b) u-channel Feynman diagrams.	159
4.10 Electron self energy as an example of closed loop diagram.	160
5.1 Interaction of a charged particle with the electromagnetic field.	169
5.2 Four point Fermi interaction.	170
5.3 (a) Effect of parity transformation on the process ${}^{60}\text{Co} \rightarrow {}^{60}\text{Ni}^* + e^- + \bar{\nu}_e$ (b) Parity transformation of a muon decaying into a positron.	179
5.4 Recoil momentum spectrum for the decay ${}^6\text{He} \rightarrow {}^6\text{Li} e^- + \bar{\nu}_e$ together with the predictions for pure A and pure T couplings.	181
5.5 Helicity of neutrino indirectly determined in Goldhaber experiment.	184
5.6 Pion decay at rest.	185
5.7 Lepton current connects $\nu_e$ to $e^-$ and $\nu_\mu$ to $\mu^-$ .	190
5.8 Muon-decay at rest.	190
5.9 A schematic representation of muon decay in case of maximum electron energy ( $\epsilon = 1$ ).	195
5.10 Parity violation in the angular distribution.	195
5.11 The higher order corrections to the muon-decay: (a) Vertex correction, (b, c) Self energy correction and (d, e) Bremsstrahlung contributions.	199
5.12 Inverse muon decay: $\nu_\mu + e^- \rightarrow \mu^- + \nu_e$ .	201
5.13 First order, second order, and higher order diagrams in $V - A$ theory for $\nu_e - e^-$ scattering.	205

5.14 $\nu_l - l^-$ scattering mediated by $W^+$ boson.	206
5.15 Feynman diagram depicting $\nu_\mu \bar{\nu}_\mu \rightarrow W^+ W^-$ via $\mu^-$ exchange.	208
5.16 $e^- e^+ (\mu^- \mu^+) \rightarrow \tau^- \tau^+$ process mediated by a photon exchange.	210
5.17 The production cross section ratio $R_{eX}^{2p} = \frac{\sigma(e^+ e^- \rightarrow eX)}{\sigma(e^+ e^- \rightarrow \mu^+ \mu^-)}$ vs. the CM energy $\sqrt{s}$ for all $eX$ events with no detected photons.	211
5.18 Normalized (left) electron and (right) muon energy distributions for tau decays compared with V - A (solid line) and V+A (dashed line) spectra.	213
6.1 Feynman diagram for the pion decay at rest, $\pi^-(p) \rightarrow l^-(k) + \bar{\nu}_l(k')$ .	219
6.2 $\beta$ -decay of $A = 12$ nuclei, viz. $^{12}\text{B}$ , $^{12}\text{N}$ , and $^{12}\text{C}^*$ .	228
6.3 Coupling of the axial vector current to a particle of zero mass.	230
6.4 Top panel: (left) pion and (right) kaon decay. Bottom panel: (left) neutron and (right) lambda decay.	238
6.5 Internal Bremsstrahlung and direct emission radiative decays.	252
6.6 Radiative decays corresponding to one quark, two quark, and three quark, transitions.	253
6.7 $K^0 \rightarrow \bar{K}^0$ oscillation via two W bosons exchange.	255
6.8 Probability of neutral kaon oscillation. (a) represents the survival probability of $K^0$ , that is, $P(K^0 \rightarrow K^0)$ , while (b) represents the oscillation probability, $P(K^0 \rightarrow \bar{K}^0)$ , for a fixed $\Delta m \neq 0$ .	256
6.9 Feynman diagrams for the $K^0$ decay into $\mu^+ \mu^-$ with a (a) $u$ -quark exchange and (b) $c$ -quark exchange.	261
6.10 Quark mixing.	261
6.11 Leptonic weak decays of mesons.	266
6.12 Semileptonic weak decays of charmed mesons.	267
6.13 Semileptonic weak decays of charmed baryons.	267
6.14 Nonleptonic decays of charmed mesons.	268
6.15 W boson exchanges.	269
6.16 QCD corrections.	269
6.17 Nucleon–nucleon scattering through a W exchange.	270
6.18 Nucleon–nucleon potential in the meson exchange model. (a) Parity-conserving (PC) potential, (b) PV potential $g\pi(\rho)NN$ and $f\pi(\rho)NN$ are the parity conserving and parity violating couplings.	271
6.19 Gluon radiative diagrams.	272
6.20 Penguin diagrams.	273
7.1 (Left) In a global transformation, $\alpha$ has a fixed value; therefore, $e^{i\alpha}$ has a constant value throughout space–time. (Right) In a local transformation, $e^{i\alpha(x)}$ , where $x = x_\mu$ is a point in space–time, has a different value at each space–time point.	281
7.2 Feynman diagrams representing different terms of Eq. (7.72).	294
7.3 (a) Quark–gluon vertex, (b) triple gluon vertex, and (c) four gluon vertex.	296

- 10.11 (a) Quasielastic and (b) Elastic  $\nu$  scattering processes on nucleon ( $N = n, p$  and  $N' = p, n$ ) target. 403
- 10.12 Kinematics in the Lab frame showing  $\nu_l + n \rightarrow l^- + p$  scattering process. 407
- 10.13 Axial mass  $M_A$  extractions from (quasi)elastic neutrino and antineutrino scattering experiments on hydrogen and deuterium targets (left) and from charged pion electroproduction experiments (right). 410
- 10.14  $\frac{d\sigma}{dq^2}$  vs.  $-q^2$  for the  $\nu_\mu + n \rightarrow \mu^- + p$  (left panel) and  $\bar{\nu}_\mu + p \rightarrow \mu^+ + n$  (right panel) at the different (anti)neutrino energies, viz.,  $E_{\nu_\mu(\bar{\nu}_\mu)} = 500$  MeV and 1 GeV and at  $M_A = 1.026$  and 1.2 GeV. 413
- 10.15  $\sigma$  vs.  $E_{\nu_l(\bar{\nu}_l)}$  for the  $\nu_l + n \rightarrow l^- + p$  (left panel) and  $\bar{\nu}_l + p \rightarrow l^+ + n$  (right panel) with  $l = e, \mu$  at  $M_A = 1.026$  and 1.2 GeV. 413
- 10.16  $\sigma$  vs.  $E_{\nu_\mu(\bar{\nu}_\mu)}$  for the process  $\nu_\mu + n \rightarrow \mu^- + p$  (left panel) and  $\bar{\nu}_\mu + p \rightarrow \mu^+ + n$  (right panel) for different combinations of  $M_A$ , and  $g_2^R(0)$ . 414
- 10.17 Feynman diagram for the process  $\bar{\nu}_l(k) + N(p) \rightarrow l^+(k') + Y(p')$ , where  $N(= p, n)$  and  $Y(= \Lambda, \Sigma^0, \Sigma^-)$  represents the initial nucleon and the final hyperon, respectively. 415
- 10.18  $\sigma$  vs.  $E_{\bar{\nu}_\mu}$ , for  $\bar{\nu}_\mu + p \rightarrow \mu^+ + \Lambda$  process. 418
- 10.19 (a) Momentum and polarization directions of the final baryon and lepton.  $\hat{e}_L^{h,l}$ ,  $\hat{e}_P^{h,l}$ , and  $\hat{e}_T^{h,l}$  represent the orthogonal unit vectors corresponding to the longitudinal, perpendicular, and transverse directions with respect to the momentum of the final hadron in (b) and the final lepton in (c). 420
- 10.20  $P_L^\Lambda(Q^2)$  vs.  $Q^2$  (left panel) and  $P_P^\Lambda(Q^2)$  vs.  $Q^2$  (right panel) for the process  $\bar{\nu}_\mu + p \rightarrow \mu^+ + \Lambda$  at the incoming antineutrino energy,  $E_{\bar{\nu}_\mu} = 1$  GeV for the polarized  $\Lambda$  in the final state, at different values of  $g_2^R(0)$ . 421
- 10.21  $P_L^\Lambda(Q^2)$  vs.  $Q^2$  (left panel),  $P_P^\Lambda(Q^2)$  vs.  $Q^2$  (middle panel) and  $P_T^\Lambda(Q^2)$  vs.  $Q^2$  (right panel) for the process  $\bar{\nu}_\mu + p \rightarrow \mu^+ + \Lambda$  at the incoming antineutrino energy,  $E_{\bar{\nu}_\mu} = 1$  GeV for the polarized  $\Lambda$  in the final state, at different values of  $g_2^I(0)$ . 422
- 11.1 Generic Feynman diagrams representing the charged and neutral current induced inelastic processes given in Table 11.1. 426
- 11.2 Generic Feynman diagrams representing the non-resonant background terms contributing to the inelastic processes. 426
- 11.3 Feynman diagram for the charged current induced spin 1/2 and 3/2 resonance production (left panel) and for the decay of the resonance into a nucleon and a pion (right panel). 431
- 11.4 Diagrammatic representation of the helicity amplitudes,  $A_{1/2}$  (top) and  $S_{1/2}$  (bottom) for spin 1/2 and 3/2 resonances. 435
- 11.5 Feynman diagram of the process  $R_{\frac{1}{2}, \frac{3}{2}} \rightarrow Nl^- \bar{\nu}_l$  through pion decay. 438
- 11.6  $Q^2$  dependence of different form factors of  $S_{11}(1650)$  resonance. 440
- 11.7  $Q^2$  dependence of different form factors of  $P_{11}(1440)$  resonance. 440
- 11.8 Feynman diagram for the decay of a spin 1/2 resonance to a nucleon and a pion. 445

- 11.9 Feynman diagram for the charged current (left) and neutral current (right) induced spin 3/2 resonance production. 447
- 11.10 Diagrammatic representation of the helicity amplitude,  $A_{3/2}$  for spin 3/2 resonances. 448
- 11.11  $Q^2$  dependence of different form factors of  $D_{13}$  resonance. From left to right panel:  $C_3^V$ ,  $C_4^V$ , and  $C_5^V$  as mentioned in Eq. (11.82);  $C_5^A$  as mentioned in Eq. (11.97); and  $C_i^{n,p}$ ,  $i = 3, 4, 5$  as mentioned in Eqs. (11.84)–(11.86). 452
- 11.12  $Q^2$  dependence of different form factors of  $P_{13}$  resonance. From left to right panel:  $C_3^V$ ,  $C_4^V$ , and  $C_5^V$  as mentioned in Eq. (11.82);  $C_5^A$  as mentioned in Eq. (11.97); and  $C_i^{n,p}$ ,  $i = 3, 4, 5$  as mentioned in Eqs. (11.84)–(11.86). 452
- 11.13 Feynman diagram for the decay of a spin 3/2 resonance in a nucleon and a pion. 455
- 11.14 Generic Feynman diagrams for the  $s$ ,  $t$ , and  $u$  channel Born terms. 464
- 11.15 Effect of explicit symmetry breaking. 470
- 11.16 Figure depicting the infinitely steep potential in the  $\sigma$  direction. 470
- 12.1 Feynman diagrams contributing for the process  $W^i N \rightarrow N' \pi^{\pm,0}$ , where ( $W^i \equiv W^\pm$ ;  $i = \pm$ ) for the charged current processes and ( $W^i \equiv Z^0$ ;  $i = 0$ ) for the neutral current processes with  $N, N' = p$  or  $n$ . 481
- 12.2 Total scattering cross section for the processes  $\nu_\mu p \rightarrow \mu^- p \pi^+$  (left panel) and  $\nu_\mu n \rightarrow \mu^- p \pi^0$  (right panel). 489
- 12.3 Total scattering cross section for neutral current neutrino induced pion production processes with deuteron effect and with no cut on  $W$ . 490
- 12.4 Feynman diagrams for the processes  $\nu/\bar{\nu}(k) + N(p) \rightarrow \mu^\mp(k') + \eta(p_\eta) + N'(p')$ . 491
- 12.5 Total scattering cross section for the charged current  $\eta$  meson production for the processes  $\nu_\mu + n \rightarrow \mu^- + \eta + p$  and  $\bar{\nu}_\mu + p \rightarrow \mu^+ + \eta + n$ . 493
- 12.6 Feynman diagrams corresponding to the (anti)neutrino induced  $\Delta S = 0$  associated particle production processes. 495
- 12.7 Cross section for neutrino (left) and antineutrino (right) induced  $|\Delta S| = 0$  associated kaon production processes. 498
- 12.8 Feynman diagrams for the process  $\nu N \rightarrow l N' K$ . The first row from left to right: contact term (labeled CT in the text), kaon pole term (KP); second row:  $u$  channel diagram ( $C\Sigma$ ,  $C\Lambda$ ) and pion (eta) in flight ( $\pi P$ ,  $\eta P$ ). 499
- 12.9 Contribution of the different terms to the total scattering cross section for the  $\nu_\mu + p \rightarrow \mu^- + K^+ + p$  (left panel) and  $\nu_\mu + n \rightarrow \mu^- + K^0 + p$  (right panel) processes. 501
- 12.10 Feynman diagrams for the process  $\bar{\nu} N \rightarrow l N' \bar{K}$ . The first row from left to right:  $s$  channel  $\Sigma, \Lambda$  propagator (labeled SC in the text),  $s$  channel  $\Sigma^*$  resonance (SCR); second row: kaon pole term (KP); contact term (CT) and last row: pion(Eta) in flight ( $\pi P/\eta P$ ). 502
- 12.11 Total scattering cross section for the processes  $\bar{\nu}_\mu p \rightarrow \mu^+ p K^-$  and  $\bar{\nu}_\mu n \rightarrow \mu^+ n K^-$ . 505
- 13.1  $e^- -^{12}\text{C}$  scattering cross section (a) Left:  $\theta = 80^\circ$ ,  $Q^2 \rightarrow 0.06 \text{ GeV}^2$ . The elastic peak is evident at low energy transfer; the excitation of nuclear levels can be seen with increase in energy transfer. (b) Right:  $\theta = 135^\circ$ ,  $Q^2 \sim 0.1 \text{ GeV}^2$ . 508

- 13.2  $e^- - ^4\text{He}$  scattering cross section for the beam energy  $E_e = 400$  MeV (a) Left:  $\theta = 45^\circ$ ,  $Q^2 \rightarrow 0.08$  GeV<sup>2</sup>, (b) Right:  $\theta = 60^\circ$ ,  $Q^2 \rightarrow 0.1$  GeV<sup>2</sup>. 509
- 13.3 Electron–proton double differential scattering cross sections at  $E_e = 4.879$  MeV; lab scattering angle  $\theta = 10^\circ$ . 510
- 13.4 (a) Left: electron–muon scattering (b) Center: electron–proton elastic scattering, and (c) Right: electron–proton deep inelastic scattering. 511
- 13.5 Charged lepton–nucleon inclusive scattering process. 512
- 13.6 Left: A virtual photon interacting with a hadron; Right: A virtual photon interacting with a point Dirac particle inside a hadron. 515
- 13.7  $\nu W_{2p}^{\text{EM}}(\nu, Q^2)$  vs.  $Q^2$  at a fixed  $x = \frac{1}{\omega}$ . Experimental data are from SLAC. 516
- 13.8 A rapidly moving hadron. 517
- 13.9 Figure on the left depicts the incoherent sum of the contributions and figure on the right represents momentum shared by the charged partons. 517
- 13.10 Parton density distribution functions using MMHT PDFs parameterization at leading order for  $2 \text{ GeV}^2 \leq Q^2 \leq 10 \text{ GeV}^2$ . 520
- 13.11 Experimental measurements for  $\frac{F_2^{\text{en}}(x)}{F_2^{\text{ep}}(x)}$  vs.  $x$  for proton and neutron targets in the case of electromagnetic interaction. 521
- 13.12 Experimental measurements for  $F_2^{\text{ep}}(x) - F_2^{\text{en}}(x)$  vs.  $x$  for proton and neutron targets in the case of electromagnetic interaction. 521
- 13.13 Left panel: Experimental verification of the Callan–Gross relation for the spin 1/2 nature of quarks. Right panel: Experimental results for the ratio of  $R_L^{\text{EM}}(x, Q^2) = \frac{F_L^{\text{EM}}(x, Q^2)}{2xF_1^{\text{EM}}(x, Q^2)}$ . 524
- 13.14 Head on collision of a quark with a virtual photon.  $\lambda$  and  $\lambda'$  denote the initial and final helicity states. 525
- 13.15 Variation of kinematic variables such as  $x$ ,  $y$ , and  $W$  is shown in the  $Q^2 - \nu$  plane. 526
- 13.16 (a) Feynman diagram for the (anti)neutrino induced deep inelastic scattering process, (b) Summation over the final hadronic state  $X$  and its amplitude square corresponds to the scattering cross section. 527
- 13.17 Feynman representation for leptonic and hadronic vertices (left and right columns) in the case of weak interaction. Feynman diagram showing the  $\nu - N$  scattering for the summed over hadronic states  $X$ . 528
- 13.18 Feynman diagram for  $W^\pm$  interactions with quarks and antiquarks. 531
- 13.19 Experimental observations for the  $(\frac{5}{18})$ th rule. 534
- 13.20 Charged current total cross section for  $\nu_l - N$  and  $\bar{\nu}_l - N$  processes. 535
- 13.21 Feynman diagram for the neutral current  $\nu_l(\bar{\nu}_l) - N$  DIS process. 536
- 13.22 Charged and neutral current (anti)neutrino differential cross sections by CHARM collaboration. 537
- 13.23 Experimental results of nucleon structure functions for (i) Top panel: electromagnetic interaction induced DIS process, (ii) Bottom panel: weak interaction induced DIS process. 540

13.24	Diagrammatic representation of (i) upper panel: the process $\gamma^*q \rightarrow qg$ and (ii) lower panel: the process $\gamma^*g \rightarrow q\bar{q}$ .	541
13.25	Diagrammatic representation of splitting functions.	542
13.26	Evolution of parton density distribution functions using MMHT PDFs parameterization from leading order to next-to-next-leading order at $Q^2 = 2 \text{ GeV}^2$ .	543
13.27	Nachtmann variable $\xi$ and slow rescaling variable $\bar{\xi}$ vs. $x$ are shown at different values of $Q^2$ .	545
13.28	Ratio of target mass corrected weak structure functions to the structure functions without TMC by using MMHT PDFs parameterization at NLO for $Q^2 = 2 \text{ GeV}^2$ .	546
13.29	Twist-4 coefficient in the renormalon approach for $F_1^{EM}(x, Q^2)$ and $F_2^{EM}(x, Q^2)$ obtained by using MMHT PDFs parameterization at NLO for $Q^2 = 2 \text{ GeV}^2$ and $10 \text{ GeV}^2$ .	547
13.30	Experimental observations for the Adler sum rule.	550
13.31	Allowed kinematical region for $\nu_l - N$ scattering in the $Q^2 - \nu$ plane for $E_\nu=3 \text{ GeV}$ (left panel) and $E_\nu=7 \text{ GeV}$ (right panel) for $Q^2 \geq 0$ .	552
13.32	$F_2^p$ structure function data on protons from SLAC and Jefferson Lab in the resonance region for $0.06 < Q^2 < 3.30 \text{ GeV}^2$ . The solid curve is a fit to deep inelastic data at $5 \text{ GeV}^2$ .	553
13.33	$W^2$ distribution of $\nu_\mu$ -water target interactions in GENIE showing the quasi-elastic scattering, the resonance interactions, and the DIS region. The $W$ distribution is further split into the three regions, KNO scaling-based model only region, PYTHIA only region, and the transition between the two regions used in the AGKY model.	554
13.34	Diagrammatic representation of resonance excitations for $l + N \rightarrow l + R$ .	555
13.35	Top left panel: Duality for the isoscalar nucleon $F_2^{eN}$ structure function. $F_2^{eN}$ vs. $\xi$ , for $Q^2 = 0.2, 0.5, 1$ and $2 \text{ GeV}^2$ , compared with several leading twist parameterizations at $Q^2 = 10 \text{ GeV}^2$ . Top right panel: Ratio $I_2^{eN}$ of the integrated $F_2^{eN}$ in the resonance region to the leading twist functions (valence and total). Bottom left panel: Duality for the isoscalar nucleon $2xF_1^{eN}$ structure function vs. $\xi$ . The results are compared with the MRST parameterization at $Q^2 = 10 \text{ GeV}^2$ , using $F_L$ and $F_2$ (dotted) and the Callan–Gross (CG) relation, $F_2 = 2xF_1$ (dot-dashed). Bottom right panel: Ratio $I_1^{eN}$ of the integrated $2xF_1^{eN}$ in the resonance region to the leading twist function.	557
13.36	Figure from (upper) Comparison of the Rein–Sehgal $F_2$ structure functions vs. $\xi$ for neutron, proton, and the isoscalar nucleon target at $Q^2 = 0.4, 1$ and $2 \text{ GeV}^2$ (left to right in each figure) with the appropriate DIS scaling functions at $Q^2 = 10 \text{ GeV}^2$ .	559
13.37	Duality for the neutrino–nucleon $F_2^{\nu N}$ , $2xF_1^{\nu N}$ and $xF_3^{\nu N}$ structure functions in the resonance region at several $Q^2$ values, indicated against their spectra. (Left) $F_2^{\nu N}$ vs. $\xi$ . The results are compared with leading twist parameterizations (valence and total) at $Q^2 = 10 \text{ GeV}^2$ . (Center) $2xF_1^{\nu N}$ vs. $\xi$ . The results are compared with the exact expression in Eq. (11.46) (dotted) and Callan–Gross relation (dot-dashed). (Right) $xF_3^{\nu N}$ vs. $\xi$ . The results are compared with several leading twist parameterizations.	559

- 14.1 Charged current induced total scattering cross section per nucleon per unit energy of the incoming particles vs. neutrino (left panel) and antineutrino (right panel) energy. 563
- 14.2 Diagrammatic representation of Pauli blocking. 567
- 14.3 Neutrino–nucleus scattering in the impulse approximation. 569
- 14.4 Neutrino–nucleus scattering with (a) final state interactions (b) exchange currents due to mesons and (c) exchange currents due to  $\Delta$ . 570
- 14.5 Nucleons in a square well potential states. 580
- 14.6 Fermi momentum  $p_F(r)$  versus  $r$  for various nuclei. 584
- 14.7 Diagrammatic representation of the particle–hole ( $p - h$ ) excitation induced by  $W(Z)$  boson in the large mass limit of IVB ( $M_{W(Z)} \rightarrow \infty$ ). 585
- 14.8 RPA effects in the  $1p - 1h$  contribution to the  $W/Z$  self-energy, where particle–hole,  $\Delta$ –hole,  $\Delta$ – $\Delta$ , etc. excitations contribute. 590
- 14.9 Ratio  $\frac{\sigma_A/N}{\sigma_{\text{free}}}$  vs.  $E_\nu$ , for neutrino (left panel) and antineutrino (right panel) induced processes in  $^{12}\text{C}$ ,  $^{40}\text{Ar}$ ,  $^{56}\text{Fe}$ , and  $^{208}\text{Pb}$ . 596
- 14.10  $\frac{d\sigma}{dp_\mu}$  vs.  $p_\mu$  for the  $\nu_\mu(\bar{\nu}_\mu)$  induced reactions on  $^{12}\text{C}$  target at  $E_\nu = 1 \text{ GeV}$ . 597
- 14.11 The fractional suppression in cross section  $\delta\sigma_{\text{model}} (= \frac{\sigma_{\text{free}} - \sigma_{\text{model}}}{\sigma_{\text{free}}})$  vs.  $E_\nu$ , where  $\sigma_{\text{free}}$  is the cross section obtained for free nucleons and  $\sigma_{\text{model}}$  is the interacting nucleon cross section in  $^{40}\text{Ar}$  obtained by using different nuclear models. 597
- 14.12  $\Delta_I = \frac{\sigma_{\nu_e(\bar{\nu}_e)} - \sigma_{\nu_\mu(\bar{\nu}_\mu)}}{\sigma_{\nu_e(\bar{\nu}_e)}}$  for neutrino (left panel) and antineutrino (right panel) induced processes in  $^{12}\text{C}$  and  $^{40}\text{Ar}$  targets. Here,  $I$  stands for the results of the cross sections obtained (i) for the free nucleon case (solid line), (ii) in the local Fermi gas model (dashed line) and (iii) for LFG with RPA effect (dashed dotted line). 598
- 14.13 Diagrams showing some typical  $2p - 2h$  contributions arising due to the  $N - N$  and  $N - \Delta$  correlations. Solid (dashed) lines denote nucleon (pion) propagators. Double lines represent  $\Delta(1232)$  propagators. Arrows pointing to the right (left) denote particle (hole) states. 599
- 14.14 Diagrams showing some typical  $2p - 2h$  contributions arising due to the meson exchange. Solid (dashed) lines denote nucleon (pion) propagators. Arrows pointing to the right (left) denote particle (hole) states. 600
- 14.15 “Quasielastic-like”  $\nu_\mu$ - $^{12}\text{C}$  cross sections measured by MiniBooNE compared to Martini et al. calculations. 601
- 14.16 MiniBooNE flux-integrated differential cross sections  $d\sigma/dQ^2$  vs.  $Q^2$  in units of  $10^{-39} \text{ cm}^2/\text{GeV}^2$  for neutrino (upper panels) and antineutrino (lower panels) CCQE-like scattering on carbon. 601
- 15.1 In a neutrino induced reaction on a nucleon target, when a pion is produced in the nuclear medium and comes out without FSI (final state interaction). 608
- 15.2 Feynman diagram considered for coherent pion production through  $\Delta - h$  excitation. 616
- 15.3 Feynman diagrams considered for neutrino induced weak coherent pion production for  $\Delta$ -resonance. 618

- 15.4 Pion production inside a nuclear target and its interaction with the nucleons in the nucleus while coming out. 621
- 15.5 A neutrino induced reaction on a nucleon target when a pion is absorbed in the nuclear medium while coming out. 621
- 15.6 Definition of impact parameter in scattering processes at high energies. 622
- 15.7  $\frac{d\sigma}{dq^2}$  and  $\frac{d\sigma}{dp_\pi}$  for the  $\nu_\mu(\bar{\nu}_\mu)$  induced charged current one  $\pi^+(\pi^-)$  process on  $^{12}\text{C}$  target at  $E_\nu = 1$  GeV. 624
- 15.8  $\sigma$  for  $\nu_\mu(\bar{\nu}_\mu)$  induced charged current incoherent  $\pi^+(\pi^-)$  production on  $^{12}\text{C}$  target. 625
- 15.9 Comparison of theoretical and event generator calculations for the differential cross sections in pion kinetic energy with the MiniBooNE  $\nu_\mu$   $\text{CH}_2$  CC  $1\pi^+$  production data. 625
- 15.10  $\sigma(E_\nu)$  vs.  $E_\nu$  for coherent  $\pi^+$  production in  $^{12}\text{C}$  626
- 15.11 (a)  $\frac{d\sigma}{dq^2}$  vs.  $-q^2$  at  $E_\nu=1$  GeV, for coherent  $\pi^+$  production in  $^{12}\text{C}$  nucleus without (dotted), with(dashed) nuclear medium effects and with nuclear medium and pion absorption effects (solid). (b)  $\frac{d\sigma}{dk_\pi}$  vs.  $k_\pi$  at  $E_\nu=1$  GeV. 627
- 15.12  $\sigma(E_{\bar{\nu}_\mu})$  vs.  $E_{\bar{\nu}_\mu}$  for  $\pi^-$  &  $\pi^0$  production in  $^{12}\text{C}$ , in the  $\Delta$  dominance model and via intermediate hyperons. 629
- 15.13 Feynman diagrams for NC photon emission. The first two diagrams are direct and crossed baryon pole terms with nucleons and resonances in the intermediate state ( $\text{N}, \Delta(1232), \text{N}^*(1440), \text{N}^*(1520)$ , etc.) The third diagram represents t channel meson ( $\pi, \rho, \omega$ ) exchange contributions. 630
- 16.1 The deep inelastic charged lepton (a) and (anti)neutrino (b) scattering processes with bound nucleons for the electromagnetic and weak interactions, respectively. 632
- 16.2 Ratio  $R(x, Q^2) = \frac{F_{2A}(x, Q^2)}{F_{2D}(x, Q^2)}$ ; ( $A =$  target nucleus) vs.  $x$  shows the nuclear medium effects in structure function. 633
- 16.3  $\sigma_L$  and  $\sigma_T$  separation using Rosenbluth technique. 637
- 16.4 The  $A$ -dependence of the nCTEQ bound proton PDFs at the scale  $Q = 10$  GeV for a range of nuclei from the free proton ( $A = 1$ ) to lead ( $A = 208$ ). 642
- 16.5 Nuclear correction factor  $R$  for the structure function  $F_2$  in neutrino and antineutrino scattering from Fe for  $Q^2 = 5$  and  $20$  GeV $^2$ . 643
- 16.6 DIS cross section ratios as a function of  $x$  for MINERvA data points and various parameterizations of  $x$ -dependent nuclear effects. 644
- 16.7 Differential scattering cross section  $\frac{1}{E_\nu} \frac{d^2\sigma}{dx dy}$  vs.  $y$ , at different  $x$  for  $\nu_\mu - ^{56}\text{Fe}$  DIS process at  $E_\nu = 65$  GeV. 644
- 16.8 Representation of neutrino self-energy. 647
- 16.9 (a) Free field fermion propagator. (b) The term that contributes to the neutrino self-energy in the lowest order. 648
- 16.10 Diagrammatic representation of the neutrino self-energy. 649
- 16.11 Diagrammatic representation of intermediate vector boson  $W$  self-energy. 650

- 16.12 Diagrammatic representation of nucleon self-energy in the nuclear medium. 651
- 16.13 Results for  $S_h(\omega, \vec{p})$  vs.  $\omega$  are shown for (a)  $p < p_F$  and (b)  $p > p_F$  in various nuclei like  $^{12}\text{C}$ ,  $^{40}\text{Ca}$ ,  $^{56}\text{Fe}$ ,  $^{120}\text{Sn}$ , and  $^{208}\text{Pb}$ . 655
- 16.14 Neutrino self-energy diagram accounting for neutrino–meson DIS (a) the bound nucleon propagator is substituted with a meson ( $\pi$  or  $\rho$ ) propagator (b) by including particle–hole ( $1p-1h$ ), delta–hole ( $1\Delta-1h$ ),  $1p1h - 1\Delta1h$ , etc. interactions. 657
- 16.15  $R(x, Q^2) = \frac{F_{iA}^{\text{WI}}(x, Q^2)}{F_{iN}^{\text{WI}}(x, Q^2)}$ ; ( $i = 2, 3$ ) vs.  $Q^2$  in  $^{56}\text{Fe}$  and  $^{208}\text{Pb}$  using the full model. 663
- 16.16 Results are shown for the weak nuclear structure function  $F_{2A}^{\text{WI}}(x, Q^2)$  vs.  $x$  at  $Q^2 = 2, 5 \text{ GeV}^2$ , in  $^{12}\text{C}$ ,  $^{56}\text{Fe}$  and  $^{208}\text{Pb}$ . 664
- 16.17  $F_{2A}^{\text{WI}}(x, Q^2)$  vs.  $Q^2$  in  $^{56}\text{Fe}$  using the full model. 665
- 16.18  $F_{2A}^{\text{WI}}(x, Q^2)$  vs.  $Q^2$  in  $^{208}\text{Pb}$  using the full model. 666
- 16.19 Results for the ratio  $R'(x, Q^2) = \frac{5}{18} \frac{F_{iA}^{\text{WI}}(x, Q^2)}{F_{iA}^{\text{EM}}(x, Q^2)}$ ; ( $i = 1, 2$ ) are obtained by using the full model at NLO in  $A = ^{12}\text{C}$ ,  $^{56}\text{Fe}$  and  $^{208}\text{Pb}$  at  $Q^2 = 5$  and  $20 \text{ GeV}^2$ . 667
- 16.20  $\frac{1}{E_\nu} \frac{d^2\sigma_A^{\text{WI}}}{dx dy}$  vs.  $y$  are shown at different values of  $x$  for the incoming beam of energy  $E = 35 \text{ GeV}$ . 668
- 17.1 Different sources of neutrinos. 669
- 17.2 Neutrino flux as a function of neutrino energy. 670
- 17.3 Proton–proton chain. 671
- 17.4 Carbon–nitrogen–oxygen cycle. 672
- 17.5 Solar neutrino spectrum in the SSM. 673
- 17.6 Primary cosmic ray flux for the proton (top line with experimental points) and alpha particles (bottom line with experimental points). 677
- 17.7 Rigidity cut off. 678
- 17.8 Primary cosmic ray flux for the proton showing knee, ankle, and GZK region. 679
- 17.9 Schematic diagram of the production of atmospheric neutrinos. 680
- 17.10 Fractional contribution of pions and kaons to the flux of muons and neutrinos. 680
- 17.11  $\mu^+$  (solid line) and  $\mu^-$  (dashed line) fluxes as a function of the muon momentum at the INO, Super-K and South Pole by integrating over all the zenith and azimuthal bins for the two different heights from the sea level, as well as at the sea level. 681
- 17.12 All-direction averaged atmospheric neutrino flux for four sites averaging over one year. KAM stands for the SK site, INO for the INO site, SPL for the South Pole, and PYH for the Pyhäsalmi mine. 682
- 17.13 Typical antineutrino spectra. 685
- 17.14 Supernova (a) neutrino and (b) antineutrino fluxes. 690
- 17.15  $\bar{\nu}_e$  energy distributions for (1)  $^{238}\text{U}$  (solid), (2)  $^{232}\text{Th}$  (dotted), and (3)  $^{40}\text{K}$  (dash-dotted) decay. 693
- 17.16 Typical sketch of neutrino beam production by accelerators. 695

17.17 Flux of neutrinos from the older generation accelerators.	697
17.18 Set up for the narrow band beam (top) and wide band beam (bottom).	698
17.19 MiniBooNE spectra for neutrino ( $\nu_\mu$ ) (left) and antineutrino ( $\bar{\nu}_\mu$ ) (right).	698
17.20 MINERvA spectra for low energy neutrino ( $\nu_\mu$ ) (left) and antineutrino ( $\bar{\nu}_\mu$ ) (right).	698
17.21 Neutrino spectrum for on-axis and different off-axis angles obtained by the T2K experiment.	700
17.22 Neutrino spectrum obtained from the various sources when at rest: $\pi$ and $\mu$ DAR (left), KDAR (center) and IsoDAR (right).	702
17.23 Schematic diagram of a muon storage ring.	704
17.24 Neutrino (solid line) energy spectrum obtained with $^{18}\text{Ne}$ boosted at $\gamma = 250$ and antineutrino (dashed line) energy spectrum obtained with $^6\text{He}$ boosted at $\gamma = 150$ .	705
17.25 Tests of fundamental physics accessible with neutrinos of different energies.	706
18.1 The top figure represents a no oscillation case where at "A", $\nu_\alpha$ s are produced along with $l_\alpha^+$ and travel a short distance, and at "B", the same number of $l_\alpha^-$ are detected through charged current interactions ( $\nu_\alpha, l_\alpha^-$ ). The bottom figure represents an oscillation case, where at "A", $\nu_\alpha$ s are produced along with $l_\alpha^+$ and travel some distance, and at "B", some of the charged leptons of flavor $l_\beta^-$ are observed which is only possible when on the way, a few $\nu_\alpha$ get converted into $\nu_\beta$ .	712
18.2 Oscillation probability vs. $\frac{\Delta m^2}{4} \frac{L}{E}$ curve for $\sin^2 2\theta = 0.83$ . (a) shows no oscillation scenario in the region $\frac{1}{\Delta m^2} \gg \frac{L}{E}$ ; (b) shows the threshold condition for oscillation at $\frac{1}{\Delta m^2} \sim \frac{L}{E}$ and (c) shows the oscillation scenario in the region $\frac{1}{\Delta m^2} \ll \frac{L}{E}$ .	715
18.3 Transition probability curve for two-flavor neutrino oscillations for different values of $L$ .	717
18.4 Transition probability curve for two-flavor neutrino oscillations for different values of $\Delta m^2$ .	717
18.5 $L^{\text{osc}}$ curves obtained using Eq. (18.30) at $E_\nu=1$ GeV and 3 GeV, for $\Delta m^2 = 2.5 \times 10^{-3} \text{eV}^2$ and $\sin^2(2\theta)=1$ .	718
18.6 Appearance probability curve in three-flavor neutrino oscillations for different values of $\delta_{CP}$ .	728
18.7 Appearance probability curve in three-flavor antineutrino oscillations for different values of $\delta_{CP}$ .	728
18.8 A three-neutrino (mass) <sup>2</sup> spectrum in which the $\nu_2 - \nu_1$ splitting $\Delta m^2$ is much smaller than the splitting $\Delta m_{\text{big}}^2$ between $\nu_3$ and the $\nu_2 - \nu_1$ pair. The later pair may be at either the bottom or the top of the spectrum.	729
18.9 Matter density variation inside the sun.	732
18.10 Matter density variation inside the earth.	733
18.11 (a) Charged current interaction (left). (b) Neutral current interaction (right).	734
18.12 Oscillation probability for $\nu$ and $\bar{\nu}$ in matter and vacuum as illustration of matter effect. Normal and inverted hierarchy, using NOvA experiment parameters, $L = 810$ km, $E = 1\text{--}6\text{GeV}$ , $\Delta m^2 = 2.5 \times 10^{-3} \text{eV}^2$ , $\theta = 8.9^\circ$ .	740

18.13 Pictorial representation of normal neutrino mass ordering and mixing for the three (left) and four (right) neutrino picture.	746
19.1 Origin of the elements through different processes.	749
19.2 Formation of the elements in the universe through the Big Bang nucleosynthesis, stellar nucleosynthesis, and the supernova nucleosynthesis by the s- and r-processes.	750
19.3 Neutrino driven wind.	751
19.4 Timeline of the Big Bang.	753
19.5 Interstellar cloud and the shock wave hitting the cloud.	757
19.6 (a) Accumulation of the matter of the interstellar cloud in a region. (b) Hydrostatic equilibrium.	758
19.7 The Hertzsprung–Russell diagram: This classifies stars according to their luminosity, spectral class, surface temperature and evolutionary stage.	759
19.8 The future of sun (a red giant).	761
19.9 Different layers of a red supergiant.	762
19.10 Planetary nebula and the white dwarf.	763
19.11 Onion like layers in the core of a massive star ( $M_{\text{star}} = 25M_{\odot}$ ). The table shows the duration of fusion taking place during the fusion of various elements.	765
19.12 Photodisintegration of iron nucleus.	766
19.13 Type Ia supernova explosion.	768
19.14 Slow neutron capture process.	769
19.15 Rapid neutron capture process.	770
19.16 Two merging neutron stars.	772
20.1 (a) $(A, Z) \rightarrow (A, Z + 2) + e^{-} + e^{-} + \bar{\nu}_e + \bar{\nu}_e$ and (b) $(A, Z) \rightarrow (A, Z + 2) + e^{-} + e^{-}$	777
20.2 $nn \rightarrow pp + e^{-} + e^{-}$	777
20.3 Flavor changing neutral current processes in higher order loop diagrams in SM.	790
20.4 Flavor changing neutral current processes in some models beyond the standard model (BSM) physics.	790
A.1 Lorentz transformation.	795
B.1 Baryon octet	809
B.2 Pseudoscalar ( $J^P = 0^{-}$ ) and vector ( $J^P = 1^{-}$ ) meson nonet.	810

# Tables

2.1 Transformations of Dirac spinor and adjoint spinors under various symmetry operations.	92
2.2 Transformations of Dirac bilinear covariants under various symmetry operations.	93
5.1 $ft$ values from super-allowed Fermi decays.	175
5.2 The columns list the parent nucleus in the transition, the initial, and final spins, and the transition type. F stands for a pure Fermi transition and GT stands for a Gamow–Teller transition.	176
5.3 Nucleon operators for the decay $n \rightarrow e^- \bar{\nu}_e$ in the relativistic and non-relativistic limits, where $\sigma_{\mu\nu} = \frac{i}{2}[\gamma_\mu, \gamma_\nu]$ , and $\psi$ s are the nuclear wave functions.	178
5.4 Values of the coefficients $A$ and $B$ appearing in Eq. (5.67).	187
5.5 Electron and muon families with their respective lepton numbers.	189
5.6 Michel parameters in $V - A$ theory and their values determined in the experiments.	197
5.7 Decay modes of tau lepton.	212
5.8 Michel parameters of the $\tau$ lepton.	214
6.1 Quarks are spin $\frac{1}{2}$ fermions with baryon number $\frac{1}{3}$ ; they have been assigned positive parity. $Q( e )$ is the quark's charge in the units of electronic charge, $I$ is the isospin and its 3rd component is $I_3$ . $S$ , $C$ , $B$ , and $T$ stand for the strangeness, charm, bottom, and top quantum numbers, respectively.	215
6.2 Mesons ( $J^P = 0^-$ ) and their properties: quark content, $J$ represents angular momentum, $P$ the parity, $I$ the isospin, $I_3$ the 3rd component of $I$ , $S$ the strangeness, $M$ the mass and $\tau$ the lifetime of a given meson.	216
6.3 Baryons ( $J^P = \frac{1}{2}^+$ ) and their properties: quark content, isospin( $I$ ) and its 3rd component ( $I_3$ ), total angular momentum ( $J$ ) and parity ( $P$ ), strangeness ( $S$ ), mass ( $M$ ), and lifetime( $\tau$ ).	216
6.4 Decay modes with the corresponding branching ratios of charged as well as neutral pions and kaons.	218

6.5 Decay modes of hyperons with branching ratios. Neutron decays to $pe^- \bar{\nu}_e$ with 100% branching ratio.	218
6.6 Production modes of kaons.	234
6.7 Cabibbo model predictions for the $\beta$ -decays of baryons. Here, for simplicity, we have dropped the superscript $A$ from $F, D$ in $g_1(0)$ and $g_1/f_1$ . The variables used in the table are defined as: $V_{ud} = \cos \theta_C$ , $V_{us} = \sin \theta_C$ , $\mu_p = 1.7928\mu_N$ , and $\mu_n = -1.9130\mu_N$ .	244
6.8 Results from $\sin \theta_C$ analysis using measured $g_1/f_1$ values.	247
6.9 Nonleptonic decays of hyperons.	251
6.10 Predicted $c\bar{c}$ and $b\bar{b}$ states with principal quantum numbers $n=1$ and 2, and radial quantum number $n_r = n - L$ , compared with experimentally observed states. Masses are given in $MeV/c^2$ .	265
6.11 Some of the observed parity violating effects in nucleons and nuclei. $A_L$ represents asymmetry and $P_L$ represents the polarization of the hadron(photon) in the final state.	271
8.1 Weak isospin ( $I$ ), its third component ( $I_3$ ), charge ( $Q( e )$ ) and hypercharge ( $Y = 2(Q - I_3)$ ) of leptons and scalar mesons in the W-S model with $l = e, \mu, \tau$ .	320
8.2 Couplings of the leptons to $Z_\mu$ field.	335
8.3 Isospin ( $I$ ), its third component ( $I_3$ ), charge ( $Q( e )$ ), and hypercharge ( $Y = 2(Q - I_3)$ ) of the first and second generation of quarks, that is, ( $u, d$ ) and ( $c, s$ ) quarks.	336
8.4 Couplings of the quarks ( $u, d$ ) to $Z_\mu$ field	337
8.5 Values of $\sin^2 \theta_W$ calculated from various data sets from different experiments.	341
8.6 Results from selected parity violating (PV) experiments. Asymmetries are given in parts per million (ppm).	343
8.7 Results from selected PV experiments. Asymmetries are given in ppm.	346
8.8 Experimental determinations of the ratios $g_1/g_v$ .	355
8.9 Branching ratios of the Higgs boson decays.	360
9.1 Values of $\alpha$ , $\beta$ , and $\gamma$ for $\nu_\mu e^-$ , $\bar{\nu}_\mu e^-$ , $\nu_e e^-$ , and $\bar{\nu}_e e^-$ scattering.	373
9.2 Measured values of the total cross section and $\sin^2 \theta_W$ from different experiments for $\nu_\mu e^-$ and $\bar{\nu}_\mu e^-$ scattering at 90% confidence level.	375
9.3 Cross section of neutrino-electron $\nu_e e$ ( $\bar{\nu}_e e$ ) scattering processes from different experiments. <sup>a</sup> Region in visible energy [1.5–3.0] MeV. <sup>b</sup> Region in visible energy [3.0–4.5] MeV.	376
9.4 Magnetic moment of neutrinos as measured by the different experiments.	377
10.1 Recent measurements of the axial dipole mass ( $M_A$ ).	410
11.1 Charged and neutral current induced inelastic processes. $N, N'$ represents protons and neutrons, $Y = \Lambda, \Sigma$ represents hyperons, $K = K^+, K^0$ represents kaons, $\bar{K} = K^-, \bar{K}^0$ represents antikaons, and $l = e, \mu, \tau$ represents leptons.	426
11.2 Properties of the spin 1/2 resonances available in the Particle Data Group (PDG) Breit-Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , parity $P$ , and the branching ratio of different meson-baryon interactions like $N\pi$ , $N\eta$ , $K\Lambda$ , and $K\Sigma$ .	427

11.3	Properties of the spin 3/2 resonances available in the PDG with Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , parity $P$ , and the branching ratio of different meson–baryon interactions like $N\pi$ , $N\eta$ , $K\Lambda$ , and $K\Sigma$ .	428
11.4	Properties of the meson (pion and kaon) resonances available in PDG with Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , spin $J$ , and parity $P$ .	428
11.5	Properties of the spin 1/2 and 3/2 hyperon resonances available in PDG with Breit–Wigner mass $M_R$ , the total decay width $\Gamma$ , isospin $I$ , spin $J$ , and parity $P$ .	429
11.6	MAID parameterization of the transition form factors for the spin 1/2 resonance on a proton target. $\mathcal{A}_\alpha(0)$ is given in units of $10^{-3} \text{ GeV}^{-\frac{1}{2}}$ and the coefficients $a_1, a_2, a_4, b_1$ in units of $\text{GeV}^{-2}, \text{GeV}^{-4}, \text{GeV}^{-8}, \text{GeV}^{-2}$ , respectively. For all fits, $a_3 = 0$ . For $S_{11}(1535)$ , $S_{31}(1620)$ , and $S_{11}(1650)$ , resonance $a_2$ and $a_4$ are taken to be 0.	436
11.7	MAID parameterization for a neutron target ( $a_{2,3,4} = 0$ ) for spin 1/2 resonances.	437
11.8	MAID parameterization of the transition form factors for the spin 3/2 resonance on a proton target. Different components of this table have the same meaning as in Table 11.6.	449
11.9	MAID parameterization for neutron target ( $a_{2,3,4} = 0$ ) for spin 3/2 resonances.	449
11.10	Expressions for $O_i^\mu(V), i = 1 - 6$ and $O_i^\mu(A), i = 1 - 8$ , where $P = p + p'$ .	465
12.1	The values of constant term ( $\mathcal{A}^i$ ) appearing in Eq. (12.6), where $i$ corresponds to the nucleon pole (NP), cross nucleon pole (CP), contact term (CT), pion pole (PP), and pion in flight (PF) terms. $f_\pi$ is the pion weak decay constant and $g_1$ is nucleon axial vector coupling.	482
12.2	Coupling constant ( $\mathcal{C}^{\mathcal{R}}$ ) for spin $\frac{1}{2}$ and spin $\frac{3}{2}$ resonances in the case of charged current induced processes. Here, $f^*$ stands for $R_{1/2} \rightarrow N\pi$ coupling which for $\Delta(1232)$ resonance is $f_{N\Delta\pi}$ and is $f_{NR_{1/2}\pi}(f_{NR_{3/2}\pi})$ for spin $\frac{1}{2}$ ( $\frac{3}{2}$ ) resonances.	485
12.3	The standard form factors for weak CC transitions of the SU(3) baryon octet.	496
12.4	Constant factors appearing in the hadronic current. The upper sign corresponds to the processes with antineutrinos and the lower sign to that with neutrinos.	496
12.5	Values of the parameters appearing in the hadronic currents.	500
12.6	The physical states of the decuplet .	503
12.7	Constant factors appearing in the hadronic current.	505
13.1	Kinematical quantities for hadron and parton.	518
14.1	Parity of vector and axial vector multipoles.	573
14.2	Binding energy, and $Q$ value of the reaction for various nuclei. The last three columns are the parameters for MHO and 2pF densities. * is dimensionless for the MHO density.	584
15.1	Coefficients used in Eq. (15.24) for the calculation of $\text{Im}\Sigma_\Delta$ as a function of energy in the case of pion nuclear scattering.	612
15.2	Coefficients used for an analytical interpolation of $C(T_\pi)$ of Eq. (15.25).	613

16.1	Some of the experiments which studied (anti)neutrino induced DIS cross section and structure functions from nuclear targets.	638
16.2	The developments in the global DGLAP analysis of nPDFs since 1998. DIS: deep inelastic scattering; DY: Drell–Yan di-lepton production; nFFs: nuclear fragmentation functions.	641
17.1	Solar neutrino detectors and their main characteristics. LS stands for liquid scintillator detectors.	676
17.2	Atmospheric neutrino detectors.	684
17.3	Reactor (short baseline) neutrino detectors.	687
17.4	Reactor (long baseline) neutrino detectors.	687
17.5	Supernova neutrino detectors.	689
17.6	Geoneutrino detectors.	694
17.7	List of accelerators.	700
17.8	Accelerator (short baseline) neutrino detectors.	701
17.9	Accelerator (long baseline) neutrino detectors.	701
18.1	The impact of matter potential in various medium.	736
18.2	$\Delta m^2$ reach of experiments using various neutrino sources. These experiments have different $L$ and $E$ range.	742
18.3	Neutrino oscillation experiments sensitive to different $\theta_{ij}$ and $\Delta m_{ij}^2$ .	742
18.4	The best fit values of the three neutrino oscillation parameters from a global fit of the current neutrino oscillation data. The values (in brackets) correspond to $m_1 < m_2 < m_3$ ( $m_3 < m_1 < m_2$ ). The definition of $\Delta m^2$ used is: $\Delta m^2 = m_3^2 - \frac{m_2^2 + m_1^2}{2}$ . Thus, $\Delta m^2 = \Delta m_{31}^2 - \frac{\Delta m_{21}^2}{2} > 0$ , if $m_1 < m_2 < m_3$ and $\Delta m^2 = \Delta m_{32}^2 + \frac{\Delta m_{21}^2}{2} < 0$ , for $m_3 < m_1 < m_2$ .	743
19.1	Different neutrino processes in supernova and proto-neutron star matter.	752
20.1	Some of the measured values of $T_{1/2}^{2\nu}$ in various nuclei.	776
20.2	$T_{1/2}^{0\nu}$ and $\langle m_{\beta\beta} \rangle$ limits (90% C.L.) from the most recent experiments on $0\nu\beta\beta$ in various nuclei.	778
20.3	Table for rare leptonic processes with lepton flavour violation (LFV).	788
20.4	Rare hadronic processes with lepton flavor violation (LFV).	789
20.5	Recent experimental limits of FCNC kaon decays.	790
20.6	Recent values of various couplings $C_i$ ( $i = V, A, S, T$ ) determined from $\beta$ decays.	793
B.1	Values of the structure constants $f_{ijk}$ and $d_{ijk}$ .	810
B.2	Values of the coefficients $a$ and $b$ given in Eqs. (B.23) and (B.24).	815