

---

# Fundamentals of Solar Radiation

---

Lucien Wald

MINES PARISTECH, FRANCE



**CRC Press**

Taylor & Francis Group

Boca Raton London New York

---

CRC Press is an imprint of the  
Taylor & Francis Group, an Informa business

---

# Contents

---

<i>List of figures</i>	<i>ix</i>
<i>List of tables</i>	<i>xix</i>
<i>Preface</i>	<i>xxi</i>
<i>Acknowledgments</i>	<i>xxix</i>
<b>1 The definition of time and different time systems</b>	<b>1</b>
1.1 <i>Reminder – geocentric and geographic coordinates – angle of incidence</i>	2
1.2 <i>Orbit of the Earth around the sun – distance between the sun and the Earth</i>	5
1.3 <i>Solar declination</i>	7
1.4 <i>Definitions of time: from year to second</i>	9
1.4.1 <i>Year – astronomical season – month – week</i>	10
1.4.2 <i>Day – hour – minute – second</i>	12
1.5 <i>Mean solar time – true solar time</i>	13
1.6 <i>Coordinated universal time and legal time</i>	15
1.6.1 <i>Coordinated universal time (UTC)</i>	15
1.6.2 <i>Legal time – civil time – time zone</i>	16
1.6.3 <i>To sum up: converting the legal time to the true solar time</i>	18
<b>2 The course of the sun over an observer on the ground</b>	<b>19</b>
2.1 <i>Position of the sun seen by an observer on the ground</i>	21
2.1.1 <i>Solar zenithal angle – solar elevation – azimuth</i>	22
2.1.2 <i>Direction of the sun in the case of an inclined plane</i>	25
2.2 <i>Solar zenithal angles at solar noon during a year</i>	26
2.3 <i>Times of sunrise and sunset</i>	27
2.4 <i>Daytime – daylength</i>	30
2.5 <i>Solar angles during equinoxes and solstices</i>	31
2.6 <i>Effective solar angles</i>	36
2.6.1 <i>Methods for computing effective angles</i>	37
2.6.2 <i>How to practically use one of these methods</i>	38
<b>3 Solar radiation incident at the top of the atmosphere</b>	<b>40</b>
3.1 <i>Radiance – irradiance – irradiation</i>	41
3.1.1 <i>Radiance</i>	41

3.1.2 Irradiance	42
3.1.3 Irradiation – radiant exposure	43
3.1.4 Converting irradiance to irradiation and vice versa	44
3.2 Solar activity – extraterrestrial radiation	45
3.3 Extraterrestrial radiation received on a horizontal plane	46
3.3.1 Typical values of hourly extraterrestrial total radiation	47
3.3.2 Daily extraterrestrial radiation	51
3.4 Extraterrestrial radiation received on an inclined plane	53
3.5 Spectral distribution of the extraterrestrial radiation	56
<b>4 Radiative transfer in the atmosphere</b>	<b>60</b>
4.1 Absorption by gases in the atmosphere	61
4.2 Scattering – overview – case of the air molecules	64
4.3 Attenuation – extinction	67
4.4 Quantities related to the attenuation	67
4.4.1 Air mass	67
4.4.2 Optical depth – optical thickness	68
4.4.3 Transmittance	69
4.4.4 Linke turbidity factor	70
4.4.5 Visibility	71
4.5 Effects of aerosols on radiation	72
4.6 Effects of clouds on radiation	76
4.7 The paths of the sun rays in the atmosphere	78
4.8 Summary of contributions of atmospheric constituents to attenuation of radiation	80
<b>5 Ground reflection</b>	<b>82</b>
5.1 Reflection factor – its spectral variations	83
5.1.1 Reflection factor	83
5.1.2 Examples of spectra of reflection factor	85
5.2 Bidirectional reflectance distribution function (BRDF)	86
5.3 Albedo	88
<b>6 Solar radiation received at ground level</b>	<b>95</b>
6.1 Components of the solar radiation at ground level	96
6.1.1 Direct, diffuse, and reflected components – global radiation	96
6.1.2 Several definitions of the direct component	98
6.1.3 Diffuse fraction – direct fraction	100
6.2 Clearness index	100
6.2.1 Definition	100
6.2.2 An example of 15-min clearness indices	102
6.2.3 An example of monthly clearness indices	104
6.3 Clear-sky index	106
6.4 The prominent role of clouds	108

6.4.1	An example of hourly radiation over the North Atlantic Ocean	108
6.4.2	Map of a multi-year average of the total solar irradiance at ground	109
6.5	<i>Sunshine duration</i>	112
6.6	<i>The most important variables for the solar radiation at ground</i>	112
6.7	<i>Decoupling the effects of clear atmosphere and clouds</i>	114
<b>7</b>	<b>Spectral distribution of the solar radiation at ground</b>	<b>117</b>
7.1	<i>Spectral distribution of the irradiance in a cloud-free atmosphere</i>	119
7.1.1	Influence of the turbidity	119
7.1.2	Spectral distribution of the direct and diffuse components	121
7.1.3	Influence of the solar zenithal angle	123
7.1.4	Typical irradiances integrated over some spectral intervals in a clear atmosphere	124
7.2	<i>Spectral distribution of the irradiance in a cloudy atmosphere</i>	126
7.2.1	Influence of the cloud optical depth	126
7.2.2	Influence of the solar zenithal angle	128
7.2.3	Typical irradiances integrated over spectral intervals in cloudy conditions	129
<b>8</b>	<b>Variability – implications for estimating radiation</b>	<b>132</b>
8.1	<i>Time and space scales</i>	133
8.2	<i>Random nature of meteorological variables – ergodicity</i>	136
8.3	<i>Fluctuations around the average – small-scale variability</i>	139
8.4	<i>Measurement – sampling</i>	144
8.5	<i>Sampling theorem – detection of the smallest details</i>	147
8.5.1	Sampling theorem	147
8.5.2	What are the smallest observable details in a series of measurements?	148
<b>9</b>	<b>Ground-based instruments for measuring solar radiation at ground</b>	<b>150</b>
9.1	<i>Common ground-based instruments</i>	152
9.1.1	Thermopiles and photodiodes – calibration and classes of instruments	152
9.1.2	Pyranometer – pyrliometer – pyranometer with a shadow ring or ball	153
9.1.3	Other instruments measuring global radiation and its diffuse component simultaneously	155
9.2	<i>Typical uncertainties of pyranometers and pyrliometers</i>	157
9.3	<i>Incomplete coverage in both space and time by stations</i>	159
9.4	<i>Calculation of daily, monthly, and yearly sums and averages</i>	162
9.4.1	Ideal case of a complete time series of measurements	162
9.4.2	Calculation of climate normals in meteorology	164
9.4.3	Case of incomplete time series	165
9.4.4	Overview of the completion of a time series	167
<b>10</b>	<b>Other means for estimating solar radiation at surface</b>	<b>169</b>
10.1	<i>Meteorological analyses and reanalyses</i>	170

10.2	<i>Images acquired by Earth observing satellites</i>	172
10.2.1	Multispectral imagery	173
10.2.2	Orbits of satellites	174
10.2.3	Overview of the methods for estimating the solar radiation	176
10.3	<i>Estimate using nearby ground stations</i>	179
10.4	<i>Usual approach to estimate components from the only knowledge of global radiation</i>	181
10.5	<i>Methods using other meteorological variables</i>	183
10.5.1	Ångström's relationship between irradiation and sunshine duration	183
10.5.2	Relationships between the total radiation and that in a given spectral range	185
10.5.3	Estimate using other meteorological variables	186
<b>11</b>	<b>Control of the plausibility of measurements</b>	<b>189</b>
11.1	<i>Objective of the control of plausibility</i>	190
11.2	<i>How to control the plausibility of measurements</i>	193
11.3	<i>Checking metadata: an overview</i>	196
11.4	<i>Checking metadata: geographic location of the station</i>	197
11.5	<i>Checking metadata: time system and time stamp</i>	198
11.5.1	Overview of the most common problems	198
11.5.2	Some graphs for the visual inspection of the time-related problems	201
11.5.3	Benefit in using the hourly extraterrestrial irradiation	202
11.6	<i>Checking metadata: checking the measured quantity and its unit</i>	203
<b>12</b>	<b>Visual and automated procedures</b>	<b>207</b>
12.1	<i>The principle of visual inspection</i>	208
12.2	<i>Visual inspection of daily measurements</i>	208
12.2.1	Daily and monthly total irradiation	209
12.2.2	An example of visual control of daily spectral values	213
12.3	<i>Visual inspection of hourly and intra-hourly values</i>	214
12.4	<i>Two-dimensional representation of hourly and intra-hourly values</i>	217
12.5	<i>Automatic procedures</i>	222
12.5.1	Hourly and intra-hourly measurements	224
12.5.2	Daily measurements	225
12.5.3	Monthly measurements	226
12.5.4	Spectral radiation measurements	228
	<i>Index</i>	231

---

# Figures

---

1.1	Definition of the angle of incidence $\theta$ . (a) The general case of an inclined plane; and (b) the case of a horizontal plane.	4
1.2	Schematic view showing the orbit of the Earth around the sun. The Earth is represented in section with the equatorial plane and its axis of rotation. The distance between the sun and the Earth is given for the two extreme points and the two points for which it is equal to 1 au, with the corresponding dates.	5
1.3	Distance between the sun and the Earth as a function of the number of the day in the year.	6
1.4	Diagram showing the cases where the difference in distance between the sun and each pole is the greatest. (a) The North Pole is the closest to the sun, while it is the South Pole in (b). The Earth is drawn in cross section with the equatorial plane and its rotation axis. The angle $\delta$ is the solar declination. The star indicates the point of latitude $45^\circ$ on the sphere.	7
1.5	Schematic view of the change in solar declination $\delta$ during a revolution of the Earth around the sun. The Earth is drawn in cross section with the equatorial plane and its rotation axis. $\delta$ is given for both solstices and equinoxes with the corresponding dates. The dashed line divides the part of the orbit where the declination is positive from that where it is negative.	8
1.6	Solar declination (in radian and degree) during the year 2019.	9
1.7	Equation of time ( $t_{TST} - t_{MST}$ ) in h (left axis) and min (right axis) as a function of the day in the year.	14
2.1	Schematic view of the course of the sun in the sky for an observer located at $P$ in the northern hemisphere.	21
2.2	Hour angle $\omega$ of the sun for an observer located at $P$ in the northern hemisphere. $\omega_{sunrise}$ is the hour angle of sunrise, and $\omega_{sunset}$ is that of sunset ( $\omega_{sunset} = -\omega_{sunrise}$ ).	22
2.3	Solar angles depicting the course of the sun in the sky for an observer located at $P$ in the northern hemisphere. $\theta_S$ is the solar zenithal angle, $\gamma_S$ is the solar elevation and $\psi_S$ is the azimuth. $N$ , $E$ , $S$ and $W$ identify the cardinal points north, east, south, and west, respectively.	22

2.4	Angle $\theta$ under which the sun is seen relative to the normal to an inclined plane. The inclination and azimuth of the plane are $\beta$ and $\alpha$ , respectively.	25
2.5	Solar zenithal angles and elevation angles at solar noon during a year at latitudes $-90^\circ$ , $-45^\circ$ , $-23.45^\circ$ , $0^\circ$ , $+23.45^\circ$ , $+45^\circ$ , and $+90^\circ$ .	26
2.6	Time at sunrise in TST, MST, and UTC during the year, at longitude $5^\circ$ at two latitudes: equator and $45^\circ$ .	28
2.7	Astronomical daytime $S_0$ as a function of the number of the day in the year at several latitudes.	30
2.8	Solar zenithal angle $\theta_S$ (a) and azimuth $\Psi_S$ (b) during days of solstices and equinoxes at equator. Curves are not drawn for hours when the sun is below the horizon ( $\theta_S > 90^\circ$ ).	32
2.9	Solar zenithal angle $\theta_S$ (a) and azimuth $\Psi_S$ (b) during days of solstices and equinoxes at latitudes $-11.725^\circ$ and $11.725^\circ$ . Curves are not drawn for hours when the sun is below the horizon ( $\theta_S > 90^\circ$ ).	32
2.10	Solar zenithal angle $\theta_S$ (a) and azimuth $\Psi_S$ (b) during days of solstices and equinoxes at latitudes $-23.45^\circ$ and $23.45^\circ$ . Curves are not drawn for hours when the sun is below the horizon ( $\theta_S > 90^\circ$ ).	33
2.11	Solar zenithal angle $\theta_S$ (a) and azimuth $\Psi_S$ (b) during days of solstices and equinoxes at latitudes $-45^\circ$ and $45^\circ$ . Curves are not drawn for hours when the sun is below the horizon ( $\theta_S > 90^\circ$ ).	33
2.12	Solar zenithal angle $\theta_S$ (a) and azimuth $\Psi_S$ (b) during days of solstices and equinoxes at latitudes $-66.55^\circ$ and $66.55^\circ$ (polar circles). Curves are not drawn for hours when the sun is below the horizon ( $\theta_S > 90^\circ$ ).	34
2.13	Solar zenithal angle $\theta_S$ during days of solstices and equinoxes at poles. Curves are not drawn for hours when the sun is below the horizon ( $\theta_S > 90^\circ$ ).	34
3.1	Definition of the radiance.	42
3.2	Annual profile of the extraterrestrial irradiance $E_{0N}$ received by a plane always normal to the sun rays. The annual average is $1361 \text{ W m}^{-2}$ .	47
3.3	Daily profiles of the extraterrestrial total irradiance and irradiation received during 1 h on an horizontal plane, at several latitudes, during the days of equinoxes (left) and solstices of June (center) and December (right).	48
3.4	Annual profile of the extraterrestrial total irradiance and irradiation received during 1 h on a horizontal plane at 12:00 TST, at various latitudes.	50
3.5	Annual profile of the extraterrestrial total irradiance and irradiation received on a horizontal plane during a day at several latitudes.	52
3.6	Map of the annual average of the extraterrestrial total irradiance received on a horizontal plane and the corresponding irradiation. Latitudes are shown on the left and longitudes at the top.	53
3.7	Annual profile of the daily mean of the extraterrestrial total irradiance at latitude $45^\circ$ received by a plane (i) horizontal, (ii) inclined at $45^\circ$ facing south, and (iii) vertical facing south.	56

3.8	Typical distribution of the extraterrestrial spectral irradiance $E_{0N}(\lambda)$ received on a plane at normal incidence between 0 and 4000 nm.	57
3.9	Typical distribution of the extraterrestrial spectral irradiance $E_{0N}(\lambda)$ received on a plane at normal incidence between 200 and 1500 nm.	58
3.10	Typical distribution of the extraterrestrial spectral irradiance $E_0(\lambda)$ received on a horizontal plane between 200 and 1500 nm for three solar zenithal angles: 0°, 30°, and 60°.	59
4.1	Typical spectral distribution of the absorption coefficient for ozone between 250 and 700 nm.	62
4.2	Example of spectral distribution of the fraction of the solar flux transmitted by a moderately dry and clean atmosphere, between 250 and 4000 nm. $\lambda$ is the wavelength. O <sub>3</sub> , ozone; O <sub>2</sub> , dioxygen; H <sub>2</sub> O, water vapor; CO <sub>2</sub> , carbon dioxide; CH <sub>4</sub> , methane.	63
4.3	Schematic view of a possible scattering pattern in the case where the scattering body (small black circle) is much smaller than the wavelength of the incident radiation shown by the solid arrow.	64
4.4	Schematic view of possible scattering patterns in the case where the size of the scattering body (small black circle) is comparable to or slightly greater than the wavelength of the incident radiation shown by the solid arrow.	65
4.5	Illustration of multiple scattering within an atmospheric column.	66
4.6	Typical monthly values of the Linke turbidity factor at several locations. (Source: SoDa Service ( <a href="http://www.soda-pro.com">www.soda-pro.com</a> )).	71
4.7	Aerosol optical depth at 550 nm every 3 h during the year 2019 at Cuzco and Nazca, estimated by the Copernicus Atmosphere Monitoring Service (CAMS). (Source: SoDa Service ( <a href="http://www.soda-pro.com">www.soda-pro.com</a> )).	73
4.8	Aerosol optical depth at 550 nm every 3 h during the year 2019 at Beijing and Mexico City estimated by the CAMS. (Source: SoDa Service ( <a href="http://www.soda-pro.com">www.soda-pro.com</a> )).	75
4.9	Aerosol optical depth at 550 nm every 3 h during the year 2019 at Carpentras and Bondville, estimated by the CAMS. (Source: SoDa Service ( <a href="http://www.soda-pro.com">www.soda-pro.com</a> )).	75
4.10	Aerosol optical depths at 550 and 1240 nm every 3 h at Carpentras in April 2019 estimated by the CAMS. (Source: SoDa Service ( <a href="http://www.soda-pro.com">www.soda-pro.com</a> )).	76
4.11	Schematic view of different paths of the downwelling solar rays. (a) Solar rays do not reach the ground. (b) Solar rays reaching the flat collector are parallel and seem to come from the direction of the sun.	79
4.12	Schematic view of the different paths of the downwelling solar rays received by a flat collector at ground, showing various contributions to the diffuse component of the radiation at ground.	79
5.1	Schematic view of the reflection on the Thartar Lake and surroundings and the contribution of the ground to the diffuse component.	83
5.2	Diagram illustrating the reflection. The incident flux arrives from the direction of illumination ( $\theta, \psi$ ). It can be reflected in several directions, including the direction ( $\theta', \psi'$ ), called here observation direction.	84

- 
- |      |   |     |
|------|---|-----|
| 5.3  | Examples of reflectance spectra between 350 and 2500 nm for concrete, limestone, wet beach sand and dry playa mud. Graphs made from spectra kindly provided by the federal agency USGS of the United States.  | 86  |
| 5.4  | Examples of reflectance spectra between 350 and 2500 nm for a few types of grass and trees. Graphs made from spectra kindly provided by the federal agency USGS.  | 87  |
| 5.5  | Maps of the parameter $f_{iso}$ describing the isotropic part of the bidirectional reflectance distribution function in Southeast Asia and Australia. Upper left: A grid every 30° in latitude and longitude is superimposed. Drawn from the data set kindly provided by Mines ParisTech of France.   | 89  |
| 5.6  | Maps of the parameter $f_{iso}$ describing the isotropic part of the bidirectional reflectance distribution function in North America. Upper left: A grid every 30° in latitude and longitude is superimposed. Drawn from the data set kindly provided by Mines ParisTech of France.  | 89  |
| 5.7  | Maps of the parameter $f_{iso}$ describing the isotropic part of the bidirectional reflectance distribution function in South America. Upper left: A grid every 30° in latitude and longitude is superimposed. Drawn from the data set kindly provided by Mines ParisTech of France.  | 90  |
| 5.8  | Maps of the parameter $f_{iso}$ describing the isotropic part of the bidirectional reflectance distribution function in Africa. Upper left: A grid every 30° in latitude and longitude is superimposed. Drawn from the data set kindly provided by Mines ParisTech of France.   | 91  |
| 5.9  | Maps of the parameter $f_{iso}$ describing the isotropic part of the bidirectional reflectance distribution function in Asia. Upper left: A grid every 30° in latitude and longitude is superimposed. Drawn from the data set kindly provided by Mines ParisTech of France.   | 92  |
| 5.10 | Maps of the parameter $f_{iso}$ describing the isotropic part of the bidirectional reflectance distribution function in Europe. Upper left: A grid every 30° in latitude and longitude is superimposed. Drawn from the data set kindly provided by Mines ParisTech of France.   | 92  |
| 6.1  | Schematic view of the direct ( $B$ ), diffuse ( $D$ ), and reflected ( $R$ ) components of the radiation received by an inclined plane at ground whose inclination is $\beta$ . $\theta_S$ is the solar zenithal angle, and $\theta$ is the angle formed by the direction of the sun and the normal to the plane.   | 97  |
| 6.2  | Daily profiles of the 15-min average of total irradiance received on a horizontal plane at ground level, for the site of latitude 50° and longitude -20°, for 4 days in 2005: 01-10, 04-10, 07-10, and 10-10. Three profiles of 15-min irradiances are plotted: the actual one, the one assuming a cloudless sky (clear sky), and the extraterrestrial one. (Source of data: HelioClim-3 database and McClear clear-sky model, available at SoDa Service (soda-pro.com).) | 103 |
| 6.3  | Daily profiles of the 15-min clearness index for the site of latitude 50° and longitude -20°, for 4 days in 2005: 01-10, 04-10, 07-10 and 10-10.  |     |

	Two profiles of clearness index are plotted: the actual one and the one assuming a cloudless sky (clear sky). (Source of data: HelioClim-3 database and McClear clear-sky model, available at SoDa Service (soda-pro.com).)	103
6.4	Yearly profile of monthly averages of total irradiance received on a horizontal plane at ground level during the year 2006 at three sites of longitude 0° and latitudes 0°, 30°, and 60°, respectively. Two profiles are plotted: the actual one and the one assuming a cloudless sky (clear sky). There are not enough data during January, November, and December at 60° to obtain a reliable average. (Source of data: HelioClim-3 database and McClear clear-sky model, available at SoDa Service (soda-pro.com).)	104
6.5	Yearly profile of monthly clearness index during the year 2006 at three sites of longitude 0° and latitudes 0°, 30°, and 60°, respectively. Two profiles are plotted: the actual one and the one assuming a cloudless sky (clear sky). There are not enough data during January, November, and December at 60° to obtain a reliable clearness index. (Source of data: HelioClim-3 database and McClear clear-sky model, available at SoDa Service (soda-pro.com).)	105
6.6	Daily profiles of the 15-min clear-sky index at the site of latitude 50° and longitude -20°, for 4 days in 2005: 01-10, 04-10, 07-10, and 10-10. The clear-sky index in cloudless (clear-sky) conditions is equal to 1 by definition. (Source of data: HelioClim-3 database and McClear clear-sky model, available at SoDa Service (soda-pro.com).)	107
6.7	Daily profiles of the hourly average of total irradiance at ground level (a) and the corresponding clearness index (b). The latitude and longitude of the site are 50° and -20°, respectively. The profiles are drawn from 2 to 5 April 2005, for the actual conditions and assuming a cloudless sky (clear sky). The extraterrestrial irradiance is also plotted on the left. (Source of data: HelioClim-3 database and McClear clear-sky model, available at SoDa Service (soda-pro.com).)	109
6.8	Map of the annual average of the extraterrestrial total irradiance received on a horizontal plane and the corresponding irradiation. Latitudes are shown on the left and longitudes at the top.	110
6.9	Multi-annual (1990–2004) average of the solar total irradiance received on a horizontal plane on the ground and the corresponding irradiation. Latitudes are shown on the left and longitudes at the top.	111
7.1	Typical spectral distributions of the solar irradiance received on horizontal surfaces located one at the top of the atmosphere and the two others at ground in clear-sky conditions respectively limpid and turbid, between 200 and 2300 nm. Solar zenithal angle is 30°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere.	119
7.2	Typical spectral distributions of the clearness index for two cloudless atmospheres respectively limpid and turbid between	

- 
- 200 and 2300 nm. Solar zenithal angle is 30°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 120
- 7.3 Typical spectral distributions of the global irradiance and its components received on a horizontal surface at ground for two cloudless atmospheres: (a) a limpid and (b) a turbid, between 200 and 2300 nm. Solar zenithal angle is 30°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 121
- 7.4 Typical spectral distributions of the direct and diffuse clearness indices for two cloudless atmospheres respectively limpid and turbid, between 200 and 2300 nm. Solar zenithal angle is 30°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 122
- 7.5 Typical spectral distributions of the solar irradiance received on a horizontal surface at ground in cloudless limpid conditions between 200 and 2300 nm for three solar zenithal angles: 0°, 30°, and 60°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 123
- 7.6 Typical spectral distributions of the clearness index  $K_T$  between 200 and 2300 nm for three solar zenithal angles: 0°, 30°, and 60°, for a cloudless limpid atmosphere. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 124
- 7.7 Typical spectral distributions of the solar irradiance received on a horizontal surface located at ground in limpid clear-sky conditions, and in cloudy conditions with cloud optical depths of 5 and 15, between 200 and 2300 nm. Solar zenithal angle is 30°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 127
- 7.8 Typical spectral distributions of the clearness index in limpid clear-sky conditions, and in cloudy conditions with cloud optical depths of 5 and 15, between 200 and 2300 nm. Solar zenithal angle is 30°. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 127
- 7.9 Typical spectral distributions of the solar irradiance received on a horizontal surface located at ground in cloudy conditions, between 200 and 2300 nm, for three solar zenithal angles: 0°, 30°, and 60°. The cloud optical depth is 5. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 128
- 7.10 Typical spectral distributions of the clearness index  $K_T$  in cloudy conditions, between 200 and 2300 nm, for three solar zenithal angles: 0°, 30°, and 60°. The cloud optical depth is 5. Results from the numerical code libRadtran simulating the radiative transfer in the atmosphere. 129
- 8.1 Schematic representation of a geographic location with two plains separated by a mountain range oriented north–south, 100 km

	wide. A moisture-laden wind blows from west to east. Symbols $P1$ and $P2$ are pyranometers measuring solar irradiation.	134
8.2	Schematic view of an infinite set of identical, small clouds, whose positions relative to each other are frozen, moving above a pyranometer at a constant speed.	139
8.3	Illustration of a dense and regular network of identical pyranometers, spaced 2 km apart over an area of $100 \times 100 \text{ km}^2$ .	142
8.4	Example of variogram of hourly irradiation as a function of the distance, expressed relative to the square of the mean of the irradiation.	143
8.5	Illustration of sampling by a spaceborne imager.	146
8.6	Illustration of the influence of the sampling step. (a) The signal to sample regularly, which is a sinusoid whose period is 20 arbitrary units. (b) The results of the sampling with a step of respectively 20, 10, and 5 arbitrary units.	148
9.1	Pyranometer and its sun shield.	154
9.2	Equipment for the simultaneous measurement of the direct component of radiation at normal incidence (pyrheliometer) and the diffuse component on a horizontal plane (pyranometer). (a) Front view and (b) profile.	155
9.3	SPN1 instrument for the simultaneous measurement of the global total radiation and its diffuse component on the horizontal plane. A cover is visible under the dome. It obscures the direction of the sun for at least one thermopile at any instant.	156
9.4	Rotating shadowband irradiometers. The arm rotates for a few seconds every minute. During the rotation, it casts a shadow on the photodiode and temporarily hides the sun.	156
10.1	Example of the ground track (solid thick line) of a near-polar orbiting satellite and swath limits (dotted lines) of a spaceborne imager having a swath width of 3000 km at equator.	174
10.2	Example of images acquired by meteorological geostationary satellites. Here, a composite image made from multispectral images obtained by the Meteosat satellite on 2019-06-21 at 12:12 UTC is shown. Copyright (2019) EUMETSAT.	176
10.3	Illustration of the spatial interpolation to estimate radiation $H(t)$ at location $P$ . Measurements $H1(t)$ , $H2(t)$ , and $H3(t)$ are acquired at measuring stations $MS1$ , $MS2$ , and $MS3$ , respectively. $d1$ , $d2$ , and $d3$ are the distances between $P$ and the stations.	180
11.1	Daily irradiation $H_{day}$ measured at the meteorological station in Addis Ababa, from 1988 to 1992. The dotted line is the extraterrestrial daily irradiation $H_{0day}$ . (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru) for $H_{day}$ and SoDa Service (soda-pro.com) for $H_{0day}$ .)	191
11.2	Daily clearness index at the meteorological station in Addis Ababa, from 1988 to 1992.	192
11.3	Superimposition of daily profiles of the hourly irradiation measured at the Quezon City station, in July and August 2018. (a) Actual measurements. (b) Actual measurements till 07-31	

---

	included, then artificial shift of +1 h. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru))	201
11.4	Daily profiles of the hourly irradiation measured at the station of Quezon City and that received at the top of the atmosphere, from 12 to 18 January 2018. Measurements have been artificially shifted by +1 h after 15 January. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru) for the measurements and SoDa Service (soda-pro.com) for the extraterrestrial irradiation.)	202
11.5	Example of checking the measured quantity and its unit based on hourly measurements performed at the station of Alice Springs, in Australia, for the time stamp 13:00. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru))	204
11.6	Example of checking the measured quantity and its unit based on 10-min measurements simulated from hourly irradiations measured at the station of Alice Springs, in Australia, for the time stamp 12:10. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru))	205
12.1	Daily irradiation measured during the year 2015 at a site at latitude $-42^\circ$ and that received in cloudless conditions according to the McClear model (soda-pro.com).	209
12.2	Diffuse component of the daily irradiation measured during the year 2015 at a site at latitude $-42^\circ$ and that received in cloudless conditions according to the McClear model (soda-pro.com).	209
12.3	Direct component of the daily irradiation measured at normal incidence during the year 2015 at a site at latitude $-42^\circ$ and that received in cloudless conditions according to the McClear model (soda-pro.com).	210
12.4	Verification of the plausibility of the measurements by superimposing the measured global irradiation and that resulting from the sum of the direct and diffuse ( $B + D$ ) components for 2015. The curves are superimposed when the measurements are consistent.	212
12.5	Daily average of irradiance in UV-A measured during 2010, at the station in Kishinev, as well as that at the top of the atmosphere and that received in cloud-free conditions according to the McClear model. (Source: World Ozone and Ultraviolet Radiation Data Centre (woudc.org) for measurements and McClear service (soda-pro.com) for extraterrestrial and clear-sky irradiations.)	213
12.6	Hourly global irradiation measured during June 2018 at the station of Sioux Falls, in the United States, as well as that at the top of the atmosphere and that received in cloud-free conditions according to the McClear model. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru) for measurements and McClear service (soda-pro.com) for extraterrestrial and clear-sky irradiations.)	215
12.7	Hourly diffuse component measured during June 2018 at the station of Sioux Falls, in the United States, as well as that at the top of the atmosphere and that received in cloud-free conditions according to the McClear model. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru) for measurements and McClear service (soda-pro.com) for extraterrestrial and clear-sky irradiations.)	216

12.8	Hourly direct component at normal incidence measured during June 2018 at the station of Sioux Falls, in the United States, as well as that received in cloud-free conditions according to the McClear model. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru) for measurements and McClear service (soda-pro.com) for clear-sky irradiations.)	217
12.9	Verification of the plausibility of the measurements by superimposing the measured global irradiation and that resulting from the sum of the direct and diffuse ( $B + D$ ) components for June 2018 at Sioux Falls, United States. The curves are superimposed when the measurements are consistent.	218
12.10	Two-dimensional representation of the hourly global irradiation measured at the Kishinev station in 2018. Black is reserved for null values. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru).)	219
12.11	Two-dimensional representation of the direct component of the hourly irradiation at normal incidence, measured at the Kishinev station in 2018. Black is reserved for null values. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru).)	220
12.12	Two-dimensional representation of the hourly global irradiation measured at the Kishinev station in 2018 after introduction of artifacts. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru).)	221
12.13	Two-dimensional representation of the hourly irradiation measured at the Bukit Kototabang station in 2015 in the spectral band [700, 2900] nm. Black is reserved for null values. (Source: World Radiation Data Centre (wrdc.mgo.rssi.ru).)	222

---

# Tables

---

1.1	Correspondence between astronomical, boreal, and austral seasons	10
2.1	Yearly average of the astronomical daytime $S_0$ at various latitudes	31
3.1	Annual average, as well as minimum and maximum of the hourly average of the extraterrestrial total irradiance when the sun is at its highest during the day (12:00 TST), during a year, at various latitudes	50
3.2	Annual averages of the extraterrestrial total irradiance received by a horizontal plane at various latitudes, as well as minima and maxima of the daily mean	52
3.3	Typical values of extraterrestrial irradiance received at normal incidence in various spectral intervals and their fraction of the total solar irradiance TSI	58
4.1	The ten genera of clouds with their corresponding altitude range and phase	77
4.2	Summary of the contributions of the various constituents of the atmosphere to the attenuation of solar radiation during its downward path to the ground	81
7.1	Typical irradiances ( $\text{W m}^{-2}$ ) received at ground at normal incidence integrated over various spectral intervals for two cloudless atmospheres: one limpid and one turbid	125
7.2	Typical values of the fraction (%) of the irradiance received at normal incidence at the top of the atmosphere (left column) and on the ground (right columns) in various spectral ranges relative to the respective total irradiance in two cloudless conditions	126
7.3	Typical irradiances ( $\text{W m}^{-2}$ ) received at ground at normal incidence integrated over various spectral intervals for two cloudy atmospheres composed of a cloud of optical depth of respectively 5 and 15	130
7.4	Typical values of the fraction (%) of the irradiance received at normal incidence at the top of the atmosphere (left column) and on the ground (right columns) in various spectral ranges relative to the respective total irradiance under cloudy conditions	131
9.1	Relative uncertainties on the values measured by a pyrheliometer for three durations of integration depending on the category of quality of the measurements, according to the World Meteorological Organization	157

9.2	Relative uncertainties on the daily values measured by a pyranometer depending on the category of quality of the measurements, according to the World Meteorological Organization	157
9.3	Relative uncertainties on the hourly values measured by a pyranometer depending on the category of quality of the measurements, according to the World Meteorological Organization	159
9.4	Number of stations that measured daily total irradiation during at least 1 year, between 1964 and 2019, according to the database of the World Meteorological Organization, in each of the seven regions	160
10.1	Correction factors $f_{clear}$ and $f_{cloud}$ for clear-sky and cloudy conditions, respectively, for each wavelength $\lambda$	187