

Introduction to Effective Field Theory

Thinking Effectively about Hierarchies of Scale

C. P. BURGESS

McMaster University and Perimeter Institute for Theoretical Physics



CAMBRIDGE
UNIVERSITY PRESS

Contents

| | |
|------------------------------|----------------|
| <i>List of Illustrations</i> | <i>page xi</i> |
| <i>List of Tables</i> | xvii |
| <i>Preface</i> | xix |
| <i>Acknowledgements</i> | xxi |

Part I Theoretical Framework 1

| | |
|--|-----------|
| 1 Decoupling and Hierarchies of Scale | 5 |
| 1.1 An Illustrative Toy Model \diamond | 6 |
| 1.1.1 Semiclassical Spectrum | 6 |
| 1.1.2 Scattering | 7 |
| 1.1.3 The Low-Energy Limit | 9 |
| 1.2 The Simplicity of the Low-Energy Limit \diamond | 9 |
| 1.2.1 Low-Energy Effective Actions | 10 |
| 1.2.2 Why It Works | 11 |
| 1.2.3 Symmetries: Linear vs Nonlinear Realization | 13 |
| 1.3 Summary | 16 |
| Exercises | 16 |
| 2 Effective Actions | 18 |
| 2.1 Generating Functionals – A Review \heartsuit | 18 |
| 2.1.1 Connected Correlations | 21 |
| 2.1.2 The 1PI (or Quantum) Action \spadesuit | 22 |
| 2.2 The High-Energy/Low-Energy Split \diamond | 26 |
| 2.2.1 Projecting onto Low-Energy States | 26 |
| 2.2.2 Generators of Low-Energy Correlations \spadesuit | 28 |
| 2.2.3 The 1LPI Action | 29 |
| 2.3 The Wilson action \diamond | 32 |
| 2.3.1 Definitions | 33 |
| 2.4 Dimensional Analysis and Scaling \diamond | 39 |
| 2.4.1 Dimensional Analysis | 39 |
| 2.4.2 Scaling | 43 |
| 2.5 Redundant Interactions \diamond | 44 |
| 2.6 Summary | 48 |
| Exercises | 49 |
| 3 Power Counting and Matching | 51 |
| 3.1 Loops, Cutoffs and the Exact RG \spadesuit | 52 |
| 3.1.1 Low-Energy Amplitudes | 53 |

| | | |
|----------|---|------------|
| 3.1.2 | Power Counting Using Cutoffs | 54 |
| 3.1.3 | The Exact Renormalization Group | 59 |
| 3.1.4 | Rationale behind Renormalization \diamond | 63 |
| 3.2 | Power Counting and Dimensional Regularization \diamond | 64 |
| 3.2.1 | EFTs in Dimensional Regularization | 65 |
| 3.2.2 | Matching vs Integrating Out | 68 |
| 3.2.3 | Power Counting Using Dimensional Regularization | 71 |
| 3.2.4 | Power Counting with Fermions | 74 |
| 3.3 | The Big Picture \diamond | 76 |
| 3.3.1 | Low-Energy Theorems | 76 |
| 3.3.2 | The Effective-Action Logic \diamond | 77 |
| 3.4 | Summary | 79 |
| | Exercises | 79 |
| 4 | Symmetries | 82 |
| 4.1 | Symmetries in Field Theory \heartsuit | 82 |
| 4.1.1 | Unbroken Continuous Symmetries | 84 |
| 4.1.2 | Spontaneous Symmetry Breaking | 87 |
| 4.2 | Linear vs Nonlinear Realizations \diamond | 90 |
| 4.2.1 | Linearly Realized Symmetries | 91 |
| 4.2.2 | Nonlinearly Realized Symmetries | 93 |
| 4.2.3 | Gauge Symmetries | 99 |
| 4.3 | Anomaly Matching \spadesuit | 105 |
| 4.3.1 | Anomalies \heartsuit | 105 |
| 4.3.2 | Anomalies and EFTs | 108 |
| 4.4 | Summary | 113 |
| | Exercises | 113 |
| 5 | Boundaries | 116 |
| 5.1 | 'Induced' Boundary Conditions | 116 |
| 5.2 | The Low-Energy Perspective | 119 |
| 5.3 | Dynamical Boundary Degrees of Freedom | 122 |
| 5.4 | Summary | 123 |
| | Exercises | 124 |
| 6 | Time-Dependent Systems | 126 |
| 6.1 | Sample Time-Dependent Backgrounds \diamond | 126 |
| 6.1.1 | View from the EFT | 128 |
| 6.2 | EFTs and Background Solutions \diamond | 129 |
| 6.2.1 | Adiabatic Equivalence of EFT and Full Evolution | 129 |
| 6.2.2 | Initial Data and Higher-Derivative Instabilities \spadesuit | 132 |
| 6.3 | Fluctuations about Evolving Backgrounds \spadesuit | 137 |
| 6.3.1 | Symmetries in an Evolving Background | 138 |
| 6.3.2 | Counting Goldstone States and Currents \spadesuit | 141 |
| 6.4 | Summary | 144 |
| | Exercises | 145 |

| | | |
|----------|--|-----|
| | Part II Relativistic Applications | 147 |
| 7 | Conceptual Issues (Relativistic Systems) | 151 |
| 7.1 | The Fermi Theory of Weak Interactions \diamond | 151 |
| 7.1.1 | Properties of the W Boson | 151 |
| 7.1.2 | Weak Decays | 153 |
| 7.2 | Quantum Electrodynamics | 155 |
| 7.2.1 | Integrating Out the Electron | 156 |
| 7.2.2 | $E \gg m_e$ and Large Logs \star | 162 |
| 7.2.3 | Muons and the Decoupling Subtraction Scheme \star | 164 |
| 7.2.4 | Gauge/Goldstone Equivalence Theorems | 167 |
| 7.3 | Photons, Gravitons and Neutrinos | 169 |
| 7.3.1 | Renormalizable Interactions \diamond | 169 |
| 7.3.2 | Strength of Non-renormalizable Interactions \diamond | 171 |
| 7.3.3 | Neutrino-Photon Interactions \star | 173 |
| 7.4 | Boundary Effects | 177 |
| 7.4.1 | Surfaces between Media | 178 |
| 7.4.2 | Casimir Energies \star | 182 |
| 7.5 | Summary | 185 |
| | Exercises | 186 |
| 8 | QCD and Chiral Perturbation Theory | 188 |
| 8.1 | Quantum Chromodynamics \star | 188 |
| 8.1.1 | Quarks and Hadrons | 188 |
| 8.1.2 | Asymptotic Freedom | 190 |
| 8.1.3 | Symmetries and Their Realizations | 192 |
| 8.2 | Chiral Perturbation Theory | 195 |
| 8.2.1 | Nonlinear Realization \diamond | 195 |
| 8.2.2 | Soft-Pion Theorems \star | 199 |
| 8.2.3 | Including Baryons | 203 |
| 8.2.4 | Loops and Logs \diamond | 205 |
| 8.3 | Summary | 208 |
| | Exercises | 209 |
| 9 | The Standard Model as an Effective Theory | 212 |
| 9.1 | Particle Content and Symmetries \diamond | 213 |
| 9.1.1 | The Lagrangian | 215 |
| 9.1.2 | Anomaly Cancellation \star | 218 |
| 9.2 | Non-renormalizable Interactions | 221 |
| 9.2.1 | Dimension-Five Interactions | 222 |
| 9.2.2 | Dimension-Six Interactions | 224 |
| 9.3 | Naturalness Issues \star | 226 |
| 9.3.1 | Technical and 't Hooft Naturalness \diamond | 226 |
| 9.3.2 | The Electroweak Hierarchy Problem | 231 |
| 9.3.3 | The Cosmological Constant Problem | 236 |
| 9.4 | Summary | 238 |
| | Exercises | 239 |

| | |
|---|-----|
| 10 General Relativity as an Effective Theory | 241 |
| 10.1 Domain of Semi-Classical Gravity \diamond | 243 |
| 10.2 Time-Dependence and Cosmology \blacklozenge | 247 |
| 10.2.1 Semiclassical Perturbation Theory | 249 |
| 10.2.2 Slow-Roll Suppression | 252 |
| 10.3 Turtles All the Way Down? \blacklozenge | 257 |
| 10.3.1 String Theory | 257 |
| 10.3.2 Extra Dimensions | 264 |
| 10.4 Summary | 269 |
| Exercises | 270 |
| | |
| Part III Nonrelativistic Applications | 273 |
| | |
| 11 Conceptual Issues (Nonrelativistic Systems) | 277 |
| 11.1 Integrating Out Antiparticles \diamond | 277 |
| 11.2 Nonrelativistic Scaling \diamond | 280 |
| 11.2.1 Spinless Fields | 280 |
| 11.2.2 Spin-Half Fields | 282 |
| 11.3 Coupling to Electromagnetic Fields \blacklozenge | 284 |
| 11.3.1 Scaling | 285 |
| 11.3.2 Power Counting | 289 |
| 11.4 Summary | 293 |
| Exercises | 294 |
| | |
| 12 Electrodynamics of Nonrelativistic Particles | 296 |
| 12.1 Schrödinger from Wilson \diamond | 296 |
| 12.1.1 Leading Electromagnetic Interactions | 296 |
| 12.1.2 Matching | 298 |
| 12.1.3 Thomson Scattering | 306 |
| 12.2 Multiple Particle Species \blacklozenge | 307 |
| 12.2.1 Atoms and the Coulomb Potential | 309 |
| 12.2.2 Dipole Approximation | 311 |
| 12.2.3 HQET | 314 |
| 12.2.4 Particle-Antiparticle Systems | 318 |
| 12.3 Neutral Systems | 326 |
| 12.3.1 Polarizability and Rayleigh Scattering | 326 |
| 12.3.2 Multipole Moments | 330 |
| 12.4 Summary | 332 |
| Exercises | 333 |
| | |
| 13 First-Quantized Methods | 335 |
| 13.1 Effective Theories for Lumps \diamond | 336 |
| 13.1.1 Collective Coordinates \heartsuit | 337 |
| 13.1.2 Nonlinearly Realized Poincaré Symmetry \blacklozenge | 340 |
| 13.1.3 Other Localized Degrees of Freedom | 344 |
| 13.2 Point-Particle EFTs | 345 |
| 13.2.1 Electromagnetic Couplings | 346 |

| | | |
|---------------------------------------|---|-----|
| 13.2.2 | Gravitational Couplings | 348 |
| 13.2.3 | Boundary Conditions I | 348 |
| 13.2.4 | Thomson Scattering Revisited | 352 |
| 13.3 | PPEFT and Central Forces \blacklozenge | 353 |
| 13.3.1 | Boundary Conditions II | 354 |
| 13.3.2 | Contact Interaction | 359 |
| 13.3.3 | Inverse-Square Potentials: Fall to the Centre | 365 |
| 13.3.4 | Nuclear Effects in Atoms | 370 |
| 13.4 | Summary | 380 |
| | Exercises | 381 |
| Part IV Many-Body Applications | | 387 |
| 14 | Goldstone Bosons Again | 391 |
| 14.1 | Magnons \diamond | 391 |
| 14.1.1 | Antiferromagnetism | 392 |
| 14.1.2 | Ferromagnetism | 397 |
| 14.1.3 | Physical Applications | 401 |
| 14.2 | Low-Energy Superconductors \blacklozenge | 403 |
| 14.2.1 | Implications of the Goldstone Mode | 404 |
| 14.2.2 | Landau–Ginzburg Theory | 410 |
| 14.3 | Phonons \blacklozenge | 413 |
| 14.3.1 | Goldstone Counting Revisited | 413 |
| 14.3.2 | Effective Action | 415 |
| 14.3.3 | Perfect Fluids | 418 |
| 14.4 | Summary | 420 |
| | Exercises | 421 |
| 15 | Degenerate Systems | 423 |
| 15.1 | Fermi Liquids \diamond | 426 |
| 15.1.1 | EFT Near a Fermi Surface | 426 |
| 15.1.2 | Irrelevance of Fermion Self-Interactions | 428 |
| 15.1.3 | Marginal Interactions | 433 |
| 15.2 | Superconductivity and Fermion Pairing \blacklozenge | 436 |
| 15.2.1 | Phonon Scaling | 436 |
| 15.2.2 | Phonon-Coulomb Competition | 441 |
| 15.3 | Quantum Hall Systems \blacklozenge | 445 |
| 15.3.1 | Hall and Ohmic Conductivity | 445 |
| 15.3.2 | Integer Quantum Hall Systems | 448 |
| 15.3.3 | Fractional Quantum Hall Systems | 452 |
| 15.4 | Summary | 457 |
| | Exercises | 458 |
| 16 | EFTs and Open Systems | 461 |
| 16.1 | Thermal Fluids | 462 |
| 16.1.1 | Statistical Framework \heartsuit | 463 |
| 16.1.2 | Evolution through Conservation | 465 |

| | | |
|---|---|------------|
| 16.2 | Open Systems | 467 |
| 16.2.1 | Density Matrices \heartsuit | 468 |
| 16.2.2 | Reduced Time Evolution \diamond | 470 |
| 16.3 | Mean Fields and Fluctuations | 472 |
| 16.3.1 | The Mean/Fluctuation Split \diamond | 473 |
| 16.3.2 | Neutrinos in Matter | 476 |
| 16.3.3 | Photons: Mean-Field Evolution \spadesuit | 481 |
| 16.3.4 | Photons: Scattering and Fluctuations \spadesuit | 489 |
| 16.3.5 | Domain of Validity of Mean-Field Theory | 494 |
| 16.4 | Late Times and Perturbation Theory \spadesuit | 495 |
| 16.4.1 | Late-Time Resummation | 496 |
| 16.4.2 | Master Equations | 500 |
| 16.5 | Summary | 507 |
| | Exercises | 508 |
| Appendix A Conventions and Units | | 514 |
| Appendix B Momentum Eigenstates and Scattering | | 529 |
| Appendix C Quantum Field Theory: A Cartoon | | 539 |
| Appendix D Further Reading | | 577 |
| | <i>References</i> | 591 |
| | <i>Index</i> | 636 |

Illustrations

- | | | |
|-----|--|--------|
| 1.1 | The potential $V(\phi_r, \phi_l)$, showing its sombrero shape and the circular line of minima at $ \phi = v$ | page 7 |
| 1.2 | The tree graphs that dominate $\tilde{\phi}_r \tilde{\phi}_l$ scattering. Solid (dotted) lines represent $\tilde{\phi}_r$ ($\tilde{\phi}_l$), and ‘crossed’ graphs are those with external lines interchanged relative to those displayed | 8 |
| 1.3 | The tree graphs that dominate the $\tilde{\phi}_r \tilde{\phi}_l$ scattering amplitude. Solid (dotted) lines represent $\tilde{\phi}_r$ and $\tilde{\phi}_l$ particles | 9 |
| 2.1 | A sampling of some leading perturbative contributions to the generating functional $Z[J]$ expressed using Eq. (2.11) as Feynman graphs. Solid lines are propagators (Δ^{-1}) and solid circles represent interactions that appear in S_{int} . 1-Particle reducible and 1PI graphs are both shown as examples at two loops and a disconnected graph is shown at four loops. The graphs shown use only quartic and cubic interactions in S_{int} | 20 |
| 2.2 | The Feynman rule for the vertex coming from the linear term, S_{int} , in the expansion of the action. The cross represents the sum $\delta S/\delta\varphi^a + J_a$ | 21 |
| 2.3 | The tree graphs that dominate the $(\partial_\mu \xi \partial^\mu \xi)^2$ (panel a) and the $(\partial_\mu \xi \partial^\mu \xi)^3$ (panels b and c) effective interactions. Solid lines represent χ propagators, while dotted lines denote external ξ fields | 32 |
| 2.4 | One-loop graphs that contribute to the $(\partial_\mu \xi \partial^\mu \xi)^2$ interaction in the Wilson and 1LPI actions using the interactions of Eqs. (1.24) and (1.25). Solid (dotted) lines represent χ (and ξ) fields. Graphs involving wave-function renormalizations of ξ are not included in this list | 35 |
| 2.5 | The tree and one-loop graphs that contribute to the $(\partial_\mu \xi \partial^\mu \xi)^2$ interaction in the 1LPI action, using Feynman rules built from the Wilson action. All dotted lines represent ξ particles, and the ‘crossed’ versions of (b) are not drawn explicitly | 38 |
| 3.1 | The graph describing the insertion of a single effective vertex with \mathcal{E} external lines and no internal lines | 55 |
| 3.2 | Graphs illustrating the two effects that occur when an internal line is contracted to a point, depending on whether or not the propagator connects distinct vertices (left two figures) or ties off a loop on a single vertex (right two figures). In both cases, a double line represents the differentiated propagator. The two options respectively correspond to the terms $[\delta S_{\text{int}}/\delta\phi(p)][\delta S_{\text{int}}/\delta\phi(-p)]$ and $\delta^2 S_{\text{int}}/\delta\phi(p)\delta\phi(-p)$ appearing in the Wilson–Polchinski relation, Eq. (3.25) of the text | 60 |
| 3.3 | One-loop graphs that contribute to the $\partial_\mu \xi \partial^\mu \xi$ kinetic term in the Wilson and 1LPI actions using the interactions of Eqs. (3.43) and (3.44). Solid (dotted) lines represent χ (and ξ) fields | 69 |

- 4.1 A sketch of energy levels in the low-energy theory relative to the high-energy scale, M , and the relative splitting, v , within a global ‘symmetry’ multiplet. Three cases are pictured: panel (a) unbroken symmetry (with unsplit multiplets); panel (b) low-energy breaking ($v \ll M$) and panel (c) high-energy breaking (with $v \gtrsim M$). Symmetries are linearly realized in cases (a) and (b) but not (c). If spontaneously broken, symmetries in case (c) are nonlinearly realized in the EFT below M . (If explicitly broken in case (c) there is little sense in which the effective theory has approximate symmetry at all) 90
- 4.2 The triangle graph that is responsible for anomalous symmetries (in four spacetime dimensions). The dot represents the operator J^μ and the external lines represent gauge bosons in the matrix element $\langle gg|J^\mu|\Omega\rangle$, where $|\Omega\rangle$ is the ground state 107
- 6.1 A sketch of the adiabatic time-evolution for the energy, $E(t)$ (solid line), of a nominally low-energy state and the energy, $M(t)$ (double line), for a representative UV state. The left panel shows level crossing where (modulo level repulsion) high- and low-energy states meet so the EFT description fails. In the right panel high-energy states evolve past a cutoff, Λ (dotted line), without level crossing (so EFT methods need not fail) 144
- 7.1 The Feynman graph responsible for the decay $\tau \rightarrow e\nu_3\bar{\nu}_1$ at leading order in unitary gauge 153
- 7.2 The tree graph that generates the Fermi Lagrangian 154
- 7.3 The Feynman graph contributing the leading contribution to photon-photon scattering in the effective theory for low-energy QED. The vertex represents either of the two dimension-eight interactions discussed in the text 159
- 7.4 The Feynman graph contributing the vacuum polarization. The circular line denotes a virtual electron loop, while the wavy lines represent external photon lines 159
- 7.5 The leading Feynman graphs in QED which generate the effective four-photon operators in the low-energy theory. Straight (wavy) lines represent electrons (photons) 161
- 7.6 Schematic of the energy scales and couplings responsible for the hierarchy of interactions among gravitons, photons and neutrinos. Here the ovals represent the collection of particles at a given energy that experience renormalizable interactions with one another. Three such circles are drawn, for energies at the electron mass, m_e , the W -boson mass, M_W , and a hypothetical scale, M_g , for whatever theory (perhaps string theory) describes gravity at very high energies 172
- 7.7 Feynman graphs giving neutrino-photon interactions in the Standard Model. Graph (a) (left panel): contributions that can be regarded as low-energy renormalizations of the tree-level weak interaction. Graph (b) (middle panel): contributions generating higher-dimension interactions when integrating out the W . Graph (c) (right panel): contributions obtained when integrating out the Z . Although not labelled explicitly,

- quarks can also contribute to the loop in panel (c). Similar graphs with more photon legs contribute to neutrino/ n -photon interactions 174
- 7.8 Feynman graphs giving neutrino/single-photon interactions within the EFT below M_{ν} . Graph (a) (left panel): loop corrections to the tree-level Fermi interaction. Graph (b) (middle panel): loop corrections to tree-level higher-dimension effective four-fermion/one-photon interactions. Graph (c) (right panel): loop-generated higher-dimension effective two-fermion/one-photon interactions. Similar graphs with more photon legs describe multiple-photon interactions 176
- 7.9 Feynman graph showing how the light-by-light scattering box diagram appears in the $2 \rightarrow 3$ neutrino-photon scattering problem. The dot represents the tree-level Fermi coupling, though C and P invariance of electromagnetic interactions imply only the vector part need be used 177
- 8.1 The Feynman graphs giving the dominant contributions to pion-pion scattering in the low-energy pion EFT. The first graph uses a vertex involving two derivatives while the second involves the pion mass, but no derivatives 201
- 9.1 An example of UV physics that can generate the dimension-five lepton-violating operator within SMEFT 222
- 9.2 Graphs contributing to the Higgs mass in the extended UV theory. Solid (or dotted) lines represent S (or Higgs) fields. Graph (a) is the one-loop graph through which a massive S particle contributes at the 1-loop level; Graph (b) is the direct contribution of the effective coupling c_2 ; the effective $\Phi^\dagger\Phi$ coupling in the low-energy Wilsonian EFT. To these are to be added all other contributions (not drawn) including one-loop Standard-Model effects. What is important is that these other effects are present in both the full theory and the low-energy EFT 229
- 9.3 A new graph that contributes to the shift in c_2 when the heavy fields are integrated out in the supersymmetric UV model. Dotted lines represent the scalar Φ field, while a solid line here represents its superpartner ψ (rather than the heavy scalar S). The double line represents the superpartner χ of the heavy scalar S . All order- M^2 terms in this graph precisely cancel those coming from the left-hand graph of Fig. 9.2 in the supersymmetric limit (in which the masses and couplings in this graph are related to those of Fig. 9.2) 234
- 10.1 Cartoon of how free string levels (labelled by $N = 0, 1, 2, \dots$ and spaced by order M_s) are split at weak coupling into a ‘fine structure’ whose size is either suppressed by a power of string coupling, $g_s M_s$, or a Kaluza–Klein compactification scale 261
- 11.1 ‘Ladder’ graphs describing multiple Coulomb interactions that are unsuppressed at low energies. Solid (dashed) lines represent Φ (A_0), propagators 290
- 12.1 The graphs used when matching the fermion-fermion-photon vertex at one-loop order. Not shown explicitly are the counter-term graphs. Graphs (b), (c) and (d) contribute wave-function renormalization contributions, though gauge invariance ensures graph (d) need not be

- evaluated explicitly in a matching calculation, since fermion charge e_q does not get renormalized. Graphs (b) and (c) do contribute nontrivially through the fermion wave-function renormalization, δZ , with graph (a) contributing the rest 301
- 12.2 ‘Ladder’ graphs describing multiple Coulomb interactions that are unsuppressed at low energies. Solid lines represent electrons (Ψ propagators), double lines represent nuclei (Φ propagators) and dashed lines represent A_0 propagators 309
- 12.3 Graphs describing multiple interactions with an external Coulomb potential, $A_0(\mathbf{k}) = Ze/k^2$. Solid lines represent Ψ propagators while dashed lines capped by an ‘ \times ’ represent insertions of the external Coulomb potential 310
- 12.4 The tree graphs whose matching determine d_s and d_v to $O(\alpha)$. All graphs are evaluated for scattering nearly at threshold, with the ones on the left evaluated in QED and the ones on the right in NRQED 320
- 12.5 Loop corrections to one-photon exchange graphs whose matching contributes to d_s and d_v at $O(\alpha^2)$. Dashed lines on the NRQED (*i.e.* right-hand) side represent ‘Coulomb’ A_0 exchange 320
- 12.6 Diagrams whose matching contributes the two-photon annihilation contributions (and imaginary parts) to d_s and d_v . 320
- 12.7 One-loop t -channel matching diagrams that contribute to d_s and d_v to $O(\alpha^2)$. Vertices and self-energy insertions marked with crosses represent terms in NRQED that are subdominant in $1/m$. Dashed and wavy lines on the right-hand (NRQED) side are, respectively, Coulomb gauge A_0 and \mathbf{A} propagators. For brevity’s sake not all of the time-orderings of the \mathbf{A} propagator are explicitly drawn 321
- 12.8 The NRQED graphs contributing to the hyperfine structure at order $m\alpha^4$ (and order $m\alpha^5$). The fat vertex in graphs (a) and (b) represents c_v , and is c_s in graphs (c). The contact interaction in (d) involves d_s and d_v . 323
- 13.1 Plot of the kink solution, $\varphi(z)/v$, as a function of $\kappa(z - z_0)$ 338
- 13.2 Sketch of the world-sheet swept out in spacetime by a one-dimensional lump (*i.e.* a string) as time evolves. The world-sheet coordinates $\sigma^a = \{\tau, \sigma\}$ label points on the world-sheet while $\chi^\mu(\sigma^a)$ describes the embedding of the world-sheet into spacetime 341
- 13.3 The relative size of scales arising when setting near-source boundary conditions to the source action: R represents an actual UV scale characterizing the size of the source; a is a (much longer) size of the external physical system; ϵ is the radius between these two where boundary conditions are imposed. The precise value of ϵ is arbitrary, subject to the condition $R \ll \epsilon \ll a$ 349
- 13.4 Graph giving the leading Thomson scattering amplitude for photon scattering by a heavy charged particle in the first-quantized formulation 353
- 13.5 Sketch of a real bulk-field profile produced by a localized source in the UV theory (solid line) superimposed on the diverging profile obtained by extrapolating towards the source from outside within the external PPEFT (dotted line). Two radii, $r = \epsilon_1$ and $r = \epsilon_2$, are shown where boundary conditions are applied using the boundary action $I_b(\epsilon)$ in the external

- EFT. The ϵ -dependence of $\mathcal{I}_b(\epsilon)$ is defined to ensure that the external profile approximates the fixed real profile, no matter what particular value of ϵ is chosen. This shows how the ϵ -dependence of the effective boundary couplings is designed to reproduce the r -dependence of the real field profile as predicted by the bulk field equations 358
- 13.6 Plot of the RG flow predicted by Eq. (13.95) for λ vs $\ln(\epsilon/\epsilon_*)$ where the RG-invariant scale ϵ_* is chosen to be the unique value of ϵ for which λ either vanishes or diverges, depending on the RG-invariant sign $\eta_* = \text{sign}(\lambda^2 - 1)$ 363
- 13.7 The RG evolution predicted by Eq. (13.119) in the complex $\hat{\lambda}/|\zeta|$ plane. The left (right) panel uses a real (imaginary) value for ζ . Arrows (shading) show direction (speed) of flow as ϵ increases. Figure taken from [350] 368
- 13.8 RG flows predicted by (13.121) for $\text{Re } \hat{\lambda}/\xi$ and $\text{Im } \hat{\lambda}/\xi$ (where $\xi = |\zeta|$) for ζ real (left panel) and ζ imaginary (right panel). Each flow defines an RG-invariant scale ϵ_* defined by $\text{Re } \hat{\lambda}(\epsilon_*) = 0$, at which point $\text{Im } \hat{\lambda}(\epsilon_*) = iy_*$ is a second RG-invariant label. ϵ_* is multiply defined when ζ is imaginary. Figure taken from [350] 369
- 14.1 A cartoon illustrating how any given spin is parallel (antiparallel) to its four nearest neighbours for ferromagnetic (antiferromagnetic) order in two dimensions 392
- 14.2 Diagram of a superconducting annulus. Dashed line marks a path deep within the annulus along which $\mathbf{A} - \nabla\phi = 0$ 408
- 15.1 A cartoon of energy levels with each line representing a state, whose energy is portrayed by its vertical position. Dots indicate which levels are populated to produce the ground state of a system of non-interacting fermions. The arrow indicates the Fermi energy 424
- 15.2 A sketch illustrating the decomposition, $\mathbf{p} = \mathbf{k} + \mathbf{l}$, of a momentum vector into a part, \mathbf{k} , on the Fermi surface plus a piece, \mathbf{l} , perpendicular to it 428
- 15.3 A sketch illustrating the allowed final momenta for 2-body scattering on a Fermi surface 431
- 15.4 A sketch of a cubic Fermi surface, illustrating two special configurations with marginal scaling. In one the sum $\frac{1}{2}(\mathbf{k}_1 + \mathbf{k}_2)$ lies on a planar part of the Fermi surface. The other special configuration arises when two regions of the Fermi surface (which in general need not be planar) are related by a ‘nesting’ vector, \mathbf{n} 433
- 15.5 Feynman graphs that renormalize the density operator $\psi^*(\mathbf{p})\psi(\mathbf{p}')$ (represented by the cross) but only in the limit where $\mathbf{p}' \rightarrow \mathbf{p}$ within the effective theory of Fermi liquids. The four-point interaction is a marginal two-body coupling, as described in the text 434
- 15.6 The Feynman graph giving the leading perturbative correction to the marginal two-body interaction strength within the low-energy theory of Fermi liquids 434
- 15.7 A typical electron-phonon interaction in which emission or absorption of a phonon of momentum $\mathbf{q} = \mathbf{p}' - \mathbf{p}$ causes a transition between two low-energy electrons near the Fermi surface 437

- 15.8 Traces of longitudinal (or Ohmic) resistivity (ρ_{xx}) and Hall resistivity (ρ_{xy}) vs applied magnetic field, with plateaux appearing in the Hall plot. The Ohmic resistivity tends to zero for fields where the Hall resistivity plateaus. Figure taken from [410] 446
- 15.9 Cartoon of semiclassical Landau motion in a magnetic field, showing how orbits in the interior do not carry charge across a sample's length while surface orbits can if they bounce repeatedly off the sample's edge. Notice that the motion is chiral inasmuch as the circulation goes around the sample in a specific direction. This is a specific mechanism for the origin of surface currents in quantum Hall systems, as are required on general grounds for the low-energy EFT by anomaly matching 452
- 16.1 A plot of neutrino mass eigenvalues, m_i^2 , (for two species of neutrinos) as a function of radius, r , within the Sun (solid lines) as well as what these masses would be in the absence of vacuum mixing: $\theta_\nu = 0$ (dashed lines). The plot falls with r for electron neutrinos, since it is proportional to the density \bar{n}_e of electrons within the Sun. Resonance occurs where the dotted lines cross. A neutrino evolving adiabatically through the resonance follows a solid line and so completely converts from one unmixed species to another. Nonadiabatic evolution has a probability P_j of jumping from one branch to the other when passing through the resonance 481

Tables

| | | |
|------|--|----------|
| 7.1 | The three generations $i = 1, 2, 3$ of fermion flavours | page 152 |
| 8.1 | Quark properties | 189 |
| 8.2 | Pion properties | 192 |
| 8.3 | Theory vs experiment for low-energy pion scattering (from [189]) | 203 |
| 12.1 | Powers of $e^{n_e} \nu^{n_\nu}$ (and of α^{n_α} when $e^2 \sim \nu \sim \alpha$) appearing in leading effective couplings | 322 |
| 12.2 | Scattering lengths for $\lambda_{\text{blue}} = 400 \text{ nm}$ and $\lambda_{\text{red}} = 600 \text{ nm}$ | 329 |
| 15.1 | A comparison of some BCS predictions with experiment | 444 |
| 15.2 | The isotope effect for various superconductors (Numbers taken from reference [409].) | 445 |
| A.1 | The signs appearing in (A.34) and (A.35) for M one of the basis (A.33) of Dirac matrices | 524 |
| C.1 | The transformation properties of common quantities under parity (P), time-reversal (T) and charge-conjugation (C) | 557 |