People of Anatolia

Past, Current and Future Research in the Biological Anthropology of Türkiye

Edited by

Benjamin Irvine, Yılmaz Selim Erdal & Lutgarde Vandeput

British Institute at Ankara Monograph 58 2024

Contents

List of figures List of tables		v vi
1.	Introduction Yılmaz Selim Erdal, Benjamin Irvine	1
2.	Sex estimation and gender in ancient Anatolian populations İsmail Özer	5
3.	Stature estimation and stature changes in Anatolia Mehmet Sağır	15
4.	Re-evaluation of the prevalence of porotic hyperostosis in Anatolian populations by applying the 'osteological paradox' approach <i>Handan Üstündağ</i>	27
5.	Biological distance relations in Anatolian populations: an anthropological review <i>Serpil Eroğlu</i>	35
6.	Intentional headshaping in Anatolia: a diachronic analysis of its typological and semantic distribution <i>Valentina D'Amico</i>	49
7.	A genomic history of Neolithisation in Anatolia: preliminary insights Ayça Aydoğan, Ezgi Altınışık, Hasan Can Gemici, Çiğdem Atakuman, Füsun Özer, Mehmet Somel	63
8.	The effects of agriculture on the human body: a case study on Anatolian Neolithic populations <i>Yılmaz Selim Erdal</i>	73
9.	Diachronic changes in the frequency of caries in Anatolian past populations <i>M. Melis Koruyucu</i>	97
10.	An isotopic perspective of diachronic changes in dietary habits and subsistence practices in Anatolia <i>Benjamin Irvine</i>	109
11.	Concluding remarks Benjamin Irvine, Yılmaz Selim Erdal	123
Index		127

List of figures

Chapter 3

3.1 Stature changes in past Anatolian populations

Chapter 4

- 4.1 Porotic hyperostosis (PH) on an individual from Kültepe. Kültepe human bones archive
- 4.2 Map of Türkiye showing the locations of the sample populations

Chapter 5

- 5.1 Fenogram showing the hierarchical clustering of Anatolian groups (adapted from Eroğlu, Erdal 2006)
- 5.2 Fenogram showing hierarchical clustering of Anatolian and adjacent groups (adapted from Eroğlu, Erdal 2006)

Chapter 6

- 6.1 Composite headshaping in a subadult from Arslantepe
- 6.2 Geographical distribution of intentional headshaping in Anatolia from the Neolithic to the present day

Chapter 7

7.1 Map showing location of sites with genomic data discussed in the text

Chapter 8

- 8.1 Map showing prehistoric sites in Anatolia examined in this study
- 8.2 Osteoarthritis rates from different prehistoric settlements in Anatolia
- 8.3 Osteoarthritis rates by different time period
- 8.4 Arm bone lengths from the Neolithic to the LC-EBA
- 8.5 Robustness values of arm bones from the Neolithic to EBA
- 8.6 Rates of medial epicondylitis in prehistoric Anatolian populations
- 8.7 Aberrant dental wear in prehistoric Anatolian populations
- 8.8 Entheseal changes on arm bones at Körtik Tepe (after Delibaş 2016)

Chapter 9

- 9.1 Caries frequencies of Anatolian populations by time period
- 9.2 Diachronic changes in the frequency of caries in Anatolian populations

Chapter 10

- 10.1 Map of all sites in Anatolia with δ^{13} C and δ^{15} N stable isotope data
- 10.2 δ^{13} C and δ^{15} N values for adults from Anatolian Neolithic to Byzantine period sample populations
- 10.3 Means and standard deviations of δ^{13} C and δ^{15} N values for adults from Anatolian Neolithic to Byzantine period grouped sample populations
- 10.4 Means and standard deviations for δ^{13} C values from Anatolian populations through time. For references and sources of the data see Table 1
- 10.5 Means and SD for δ^{15} N values from Anatolian populations through time. For references and sources of the data see Table 1
- 10.6 Schematic representation of the 'hourglass' theory for changes in agriculture over time in Anatolia

List of tables

Chapter 2

- 2.1 Terminology and symbols used to determine sex from the human skeleton
- 2.2 Selected studies determining sex utilising discriminant function analysis
- 2.3 Sex determination utilising discriminant function analysis for selected parts of the adult human skeleton
- 2.4 Results of a perception survey about gender roles in modern Turkish society (following Vatandaş 2007)

Chapter 3

- 3.1 Martin's and Vandervael's stature classifications (Martin 1914; Vandervael 1943)
- 3.2 Regression formulas established by Pearson (1899) for estimating stature of females and males
- 3.3 Stature regression formulas established by Dupertuis and Hadden (1951)
- 3.4 Stature regression formulas and standard errors for white and black North Americans (Trotter, Gleser 1952)
- 3.5 Regression formulas created by Trotter and Gleser for white and black North American and Mongolian and Mexican males (Trotter, Gleser 1958)
- 3.6 Stature regression formulas created by Genoves (1967)
- 3.7 Regression formulas created by Sağır (1994) using X-Ray films
- 3.8 Regression formulas created by Şam (1994)
- 3.9 Regression formulas created by Günay (1995)
- 3.10 Stature regression formulas created for males and females (Sağır 2000)

Chapter 4

- 4.1 Percentages (%) of individuals by age group in the sample populations. Foetal (<birth), infant (birth-2.9 years), child (3-11.9 years), adolescent (12-19.9 years), young adult (20-34.9 years), middle adult (35-49.9 years), old adult (50+ years) and adult (>19 years)
- 4.2 Prevalence of porotic hyperostosis and subadult mortality rates in the sample populations

Chapter 6

6.1 Chronological and typological distribution of headshaping in Anatolia

Chapter 8

- 8.1 Maximal Length of the upper extremities in Anatolian populations from the PPNA to LC-EBA
- 8.2 Circumferences of upper extremities in Anatolian populations from the PPNA to LC-EBA
- 8.3 Frequency of medial epicondylitis in prehistoric Anatolia populations
- 8.4 Changes in the frequency of medial epicondylitis
- 8.5 Frequencies of abrasion on the anterior teeth of Anatolian populations
- 8.6 Comparative results of the different skeletal and dental aspect in foragers and agrarian peoples of prehistoric Anatolia
- 8.7 Sex differences in selected skeletal characteristics for foragers and agrarian peoples of prehistoric Anatolia

Chapter 9

9.1 Caries frequencies and dental wear scores of Anatolian populations

Chapter 10

- 10.1 Summary overview of sites with stable isotope data (δ^{13} C and δ^{15} N) used in this study. N.B. Sites that only have population means reported are notified with an *. Time periods refer to local regional chronologies used by the researchers, where possible absolute dates have been included. PPN = Pre-Pottery Neolithic; LN = Late Neolithic; ECh = Early Chalcolithic; Ch = Chalcolithic; LCh = Late Chalcolithic; EBA = Early Bronze Age; MBA = Middle Bronze Age
- 10.2 Means, standard deviations (SD) and ranges for δ^{13} C and δ^{15} N data from the Neolithic, Chalcolithic, EBA and Classical–Mid. Byzantine periods of Anatolia

- 10.3 Pairwise comparison *p*-values for δ^{13} C between the chronologically grouped sample populations. Bold indicates statistically significant
- 10.4 Pairwise comparison *p*-values for δ^{15} N between the chronologically grouped sample populations. Bold indicates statistically significant